

Research Journal of Pharmaceutical, Biological and Chemical Sciences

Evaluation of Anti-inflammatory Activity of Aqueous Leaf Extracts of *Tecoma stans* on Carrageenan Induced Paw Oedema in Rats by Using Digital Plythesmometer.

Kusuma Priya N*, Sirisha K, Sai Prathyusha TV, and Sathish Kumar M.

Department of Pharmacology, Chalapathi Institute of Pharmaceutical Sciences, Guntur, Andhra Pradesh, India.

ABSTRACT

The study of was undertaken to evaluate the anti-inflammatory activity of *Tecoma stans* leaf extracts using carrageenan induced paw oedema in rats. Paw oedema was produced by the injection of carrageenan underneath the plantar region of hind limb in rats. The study comprised of three treatment groups namely: control, standard and test (*Tecoma stans* aqueous leaf extract) all with four animals in each group. Severity of inflammation was observed and measured by using digital plythesmometer. At the end of study the test group showed significant anti-inflammatory activity, showing an overall p value <0.0001 when compared with other treatment groups i.e., control and standard, and showed p value <0.001 when compared with other treatment groups i.e. test vs standard.

Keywords: *Tecoma stans*, anti-inflammatory activity, carrageenan, paw oedema, digital plythesmometer, phlogistic agent (irritant).

Corresponding author

INTRODUCTION

Inflammation is a tissue reaction to infection, irritation or foreign substances. It is a part of the host defence mechanism [1]. Inflammation is the complex biological response of vascular tissues to harmful stimuli including pathogens, irritants, or damaged cells. It is a protective attempt by the organisms to remove the injurious stimuli as well as initiate the healing process for the tissue.

The process of inflammation is necessary in healing of wounds. Inflammation however if runs unchecked, lead to onset of disease like vasomotor rhinorrhoea, rheumatoid arthritis and atherosclerosis [2]. There are several tissue factors or mechanisms that known to be involved in the inflammatory reactions such as release of histamine, that is synthesised from the amino acid histidine by the action of histidine decarboxylase [3], bradykinin and prostaglandins [4].

The inflammatory reaction is readily produced in rats in the form of paw oedema (underneath the plantar region) with the help of irritants. Irritants generally known as phlogistic agents, some of them are carrageenan (sulphated polysaccharide 1% w/v), formalin (0.1ml of 1%), bradykinin (0.1ml of 0.005%), histamine, serotonin, mustard (0.1ml of 2.5% mustard powder suspension) or egg white (0.05ml of undiluted fresh egg white) brewer's yeast (0.1ml of 2.5%), formaldehyde, dextran (0.1ml of 1-3% of dextran solution), kaolin (0.1ml of 5% suspension) [5].

Carrageenan Induced Paw Oedema

Carrageenan which is a phlogistic agent known to produce inflammation by releasing histamine, serotonin and bradykinin and prostaglandin. Carrageenan-induced oedema is used to study the effect of drugs on acute phase of inflammation [4]. Carrageenan is a polysaccharide obtained from seaweed (Rhodophyceae). Carrageenan induced paw oedema model is known to be sensitive to cyclooxygenase inhibitors and has been used to evaluate the effect of non-steroidal anti-inflammatory agents which primarily inhibit the enzyme cyclooxygenase involved in prostaglandin synthesis [6].

Experimental Animals

Wistar rats of either sex (180-250gms) were maintained for 7 days in the animal house of Chalapathi Institute of Pharmaceutical Sciences, Guntur under standard conditions temperature (24±10C), relative humidity (45-55%) and 12:12 light: dark cycle. The animals were fed with standard rat pellet and water ad libitum. The animals were allowed to acclimatize to laboratory conditions 48h before the start of the experiment 4 rats/group was used. The experiment conducted after obtaining permission from the Institutional Animal Ethics Committee (IAEC) of Chalapathi Institute of Pharmaceutical Sciences, Guntur.

Selection of Dose and Treatment

Equipment

Digital plythesmometer:

The digital plythesmometer is an advanced instrument for screening anti-inflammatory drugs. It is designed for accurate measurement of rat and mouse paw oedema. The advanced instrument design records small differences in fluid level due to displacement when the paw is immersed. The digital read-out shows the exact volume of the paw.

Animals

Wistar albino rats of either sex (body weight: 180-250gms).

Dose

The anti-inflammatory activity of the leaf extracts of *Tecoma stans* was investigated using the carrageenan induced paw oedema. The test animals were randomly chosen and divided into three groups having four rats in each as follows:

Group-1: Control group (prepared 1%w/v solution of carrageenan and injected 0.1ml underneath the plantar region).

Group-2: Standard (Indomethacin dose 20mg/kg s.c., stock solution containing 4mg/ml of the drug and injected 0.5ml/100gms of body weight of the animal.)

Group-3: test (*Tecoma stans* leaf extract 60mg/kg)

Drug profile

Indomethacin comes under the classification of anti-inflammatory drugs as NSAID'S. Acetic acid derivatives group include indomethacin, sulindac, and etodolac [7].

Statistical Analysis

The values are expressed as mean ± SEM. The results were analysed for statistical significance using two way ANOVA followed by Dunnett's multiple comparison test.

RESULTS AND DISCUSSION

Carrageenan-induced rat paw oedema model is a suitable test for evaluating anti-inflammatory drugs, which has frequently been used to assess the anti oedematous effect of the drug [8]. Carrageenan is a strong chemical use for the release of inflammatory and pro-inflammatory mediators (prostaglandins, leukotriene, histamine, and bradykinin). Micro-organisms can be used for assessment of the anti-inflammatory property in carrageenan induced paw oedema in rats [9, 10]. The present results suggest that *Tecoma stans* leaf extract suppresses the first phase of carrageenan-induced paw oedema, thus confirming an NSAID-like property. The present study showed that *Tecoma stans* have anti-inflammatory property.




Figure 1: Anti inflammatory activity of aqueous leaf extract of *Tecoma stans* compared to the control group

Table: 1: Anti-inflammatory activity of aqueous leaf extract of *Tecoma stans*.

S.No	Treatment	Paw oedema volume(ml) measured by using digital plythesmometer			
		Mean ± SEM			
		0 min	15 min	30 min	60 min
1	Control	0.12±0.011	0.26±0.018	0.43±0.023	0.66±0.046
2	Standard (Indomethacin 20mg/kg)	0.12±0.010	0.135±0.006***	0.13±0.009***	0.12±0.011***
3	Test (60 mg/kg)	0.10±0.004	0.15±0.009	0.21±0.010*	0.36±0.018***

*p<0.05 significant value, ***p<0.001 significant value

CONCLUSION

The leaf extract of *Tecoma stans* showed anti-inflammatory activity when compared with the other treatment groups and therefore has the potential of being used in the treatment of inflammation.



REFERENCES

- [1] Kulkarni SK, Arzi Kaul PN. Japan J Pharmacol 1980;163-166.
- [2] Uma Bhandari, Vinay Kumar, Rahila Ahmad Pathan. 2010-2011;
- [3] Barar FSK. Essentials of Pharmacotherapeutics.4th Edition 2007.Pg.no 517.
- [4] Wills AL, P Montagazza and E W Horton, Eds., pp. 31–48, Academic Press, London, UK, 1969.
- [5] Gerhard Vogel H. 2nd edition pg.No760.
- [6] Sanjay B Kasture. 2006;75-76.
- [7] Lippincott's Illustrated Reviews, Pharmacology, 4th Edition.Pg.no 507.
- [8] The role of central mechanisms in the anti-inflammatory effect of amitriptyline on carrageenan-induced paws edema in rats.
- [9] Sarika Amdekar. Int J Inflam. 2012.
- [10] R Smith. Res J Pharm Biol Chem Sci 2014; 5(1):948-952.