

Research Journal of Pharmaceutical, Biological and Chemical Sciences

Concentration of N, P and K in potato tubers organically fertilized which sprinkled with Zn, Mn at different stage growth.

Jawad T M AL Fadlly*.

Agriculture College, Baghdad University, Iraq.

ABSTRACT

A field experiment of spring season 2012 was carried out in horticulture dept – college of Agriculture – university of Baghdad, to study the response of organically fertilized potatoes to foliar fertilization of Zn at 60 mg l^{-1} and Mn at 30 mg l^{-1} separately and together at ($60 \text{ mg l}^{-1} + \text{Mn at } 30 \text{ mg l}^{-1}$) at three stages of growth: vegetative growth, tuber initiation and tuber bulking and interaction between them, RCBD was used at three replication, Results showed that sprinkling with (Zn + Mn) have given higher concentration of N, P, K in tubers at 1.40, 0.28 and 2.24% respectively, Highest values of nitrogen was 1.13% and potassium was 0.235% when both nutrient were sprinkled at vegetative stage, While the highest concentration of phosphorous 0.270% obtained at tuber bulking stage when sprinkled both nutrient, treatment (both nutrient sprinkling on vegetative stage) gave higher concentration at 1.48, 0.33, 2.40% of N, P and K in tuber respectively.

Keywords: Potato, Zinc, Manganese, Organic, Foliar application.

**Corresponding author*

INTRODUCTION

Potato (*Solanum tuberosum* L) is one of the most important in the world according to the data of food and Agriculture Organization of United Nations (Flis et al., 2012), Micronutrient elements play a critical role in plants lead to increase of leaf area index and thereby increased light absorption and increase the amount of dry matter accumulation and economic yield (Ravi et al., 2008). Foliar feeding minimizes environmental pollution and improves nutrient utilization through reducing the amounts of fertilizers added to the soil (Abou-El-nour, 2002), Foliar fertilization not only improves plant growth traits, crop yields and nutrients uptake by crops (Maitlo et al., 2006) but also enhances nutrient use efficiency of crops (Fageria et al., 2009). Zinc (Zn) is known to have an important role either as metal component of enzymes or as functional, structural or regulatory cofactor of a large number of enzymes (Grotz and Guerinot, 2006), Zinc has an important role in the production of biomass furthermore, Zinc may be required for chlorophyll production, pollen function, fertilization and germination (Kaya and Higgs, 2002 and Pandey et al., 2006). Manganese is one of the main micronutrient, which has an important role in plants, low levels are absolutely necessary for normal nutrition and development of plants (Clarkson, 1988 and Migocka and Klobus, 2007), Manganese has an important role on activates several enzymes which involve to oxidation reaction, carboxylation, carbohydrates metabolism, phosphorus reaction and citric acid cycle (Jackson et al., 1978; Mukhopadhyay and Sharma, 1991 and Millaleo et al., 2010). Several researches indicated positive influence of micronutrient (Zn + Mn) application increase of yield and quantitative parameters of crops (Mousavi et al., 2007) on potato. In most of the Iraqi soils pH is high, in this type of soils solubility of microelements is less and cause decline absorbency these elements and finally requirement of plants to these elements is increasing (Uygur and Rimmer, 2000). Mohamadi (2002) found that application of Zn along with Mn to from foliar application caused increase in efficiency and quality of potato crop, Al – Fadlly (2016) reported that using Fe and Zn affected of weight of potato tuber, tuber yield of plants and total tuber yield.

Therefore, the present aims to evaluate the efficiency of using foliar application of Zinc and manganese and the combination in three stage growth on concentration of NPK in tuber potatoes.

MATERIALS AND METHODS

The study area is located in Department of Horticulture, College of Agriculture, Abu – Grabs University of Baghdad – Iraq in spring season 2012 on clay loam soil. The physical and chemical analysis of the experimental soil are shown in table 1, The experiment included 12 treatments with two factors, Factor one is control spraying water only (T_0), spraying 60 mg Zn l^{-1} (T_1), spraying 30 mg Mn l^{-1} (T_2) and spraying mixture Zn + Mn (60 + 30) mg l^{-1} , Factor two is sprayed at three stages of potato growth which were vegetative growth (F_1), tuber initiation (F_2) and tuber bulking (F_3). The experiment was laid out in factorial randomized block design with three replications, Field was divided into three blocks, each block was divided to 12 experimental plots, The area of each experimental plot was 6.75 m^2 (3 ridges 0.75 m width and 3 m in length), left a distance 1 m between blocks and experimental plots. Compost of mixture equal amount of waste cow, sheep and poultry with 50 Ton L^{-1} (Al-Fadlly, 2011) which decomposition add to all experimental units, Specification set out in table 2. Compost adds to each ridge before buried tuber 10 days silt in top of ridges with 30 cm depth 20 cm width and covered with soil. In 18 January 2012 tuber class desire were sown in top of ridges deeply at 10 – 12 cm, 25 cm the distance between tubers (Muharem and Abdul, 1987). F_1 treatments sprayed in vegetative stage at 9 April 2012, In 20 April sprayed F_2 treatments at initiation stage and F_3 treatment sprayed at tuber bulking stage in 1 May 2012. At maturity stage on 26 May 2012 tubers were harvested. Five plants from each net plot were tagged to record data, the content of nitrogen (N), phosphorus (P) and potassium (K) in dry mass of potato tubers were determined, Total nitrogen was measured by the Kjeldahl method (Bremner and Mulvaney, 1982). Total phosphorus was determined calorimetrically using ascorbic acid method described by (Watanabe and Olsen, 1965) and concentration of potassium (K) was determined in digested material using flame photometers as described by (Eppendorf and Hing, 1970). Data were analyzed statistically through the analysis of variance (ANOVA) using SAS system (SAS, 2001) and comparison among the average was calculated using the LSD test at significance level of 0.05.

Table 1: Some physical and chemical properties of soil of field experiment

Property		Value	Unit	Ref
pH		7.59	-	Richards, 1954
Ec (1:1)		3.15	dSm ⁻¹	
Particle Size	Sand	171.50	g, kg ⁻¹ soil	
	Silt	512.64		
	Clay	315.86		
Texture		Silt Clay Loam		
Bulk density		1.36	g. cm ⁻³	Richards, 1954
Gypsum		5.24	g kg ⁻¹ soil	
CEC		26.80	C mol ⁺ kg ⁻¹ soil	FAO, 2007
SOM		18.20	g kg ⁻¹ soil	Page <i>et al.</i> , 1982
Carbonate minerals		182.52		FAO, 2007
Soluble cat ion	Ca ⁺²	8.40	C mol ⁺ kg ⁻¹ soil	Page <i>et al.</i> , 1982
	Mg ⁺²	5.13		
	K ⁺¹	0.55		
	Na ⁺¹	3.95		
A available Nutrient Element	N	36.00	mg. Kg ⁻¹ soil	
	P	11.35		
	K	161.64		
	Zn	1.73		
	Mn	2.19		

Table 2: Chemical analyses of organic manures used

Parameter	Value	Unite
pH (1:5)	6.3	-
Ec (1:5)	30.37	ds m ⁻¹
C/N ratio	16.33	-
Organic C	325	g kg ⁻¹
Organic N	19.90	
Organic P	11.99	
Organic K	17.46	
Zn	220.16	ppm
Mn	174..34	
Ca	22.00	Meg L ⁻¹
Mg	11.00	
Organic matter	50.32	%
Humic	1.43	
Volvic	0.188	
Human	6.98	

RESULTS AND DISCUSSION

Nitrogen concentration in tubers

Statistical analysis results table 3. Showed the significant effect of foliar application of Zn and Mn separately and (Zn + Mn) together on the concentration of N in tubers, where T₃ foliar application (Zn + Mn) was superior on all other foliar treatment, also it gave the highest concentration of N at 1.4% in an increase of 64.71% as compared to control T₀ (water only foliar application) that was 0.85% at an increase of 30.84% when compared to T₁ (Zn foliar application) at 1.07% and increase of 20.68% compared to T₂ (Mn foliar application) at 1.16% N concentration. T₂ was superior on T₁ in N concentration at 8.41% increase. Time of application has no significant effect on N concentration in tubers, while interaction of Zn + Mn and time of

application was a significant impact in increasing N concentration in tubers, Where T₃F₁ (application of Zn + Mn at vegetative growth stage) was superior on all other interaction treatments and gave the highest N concentration in tubers at 1.48% and increase of 85% when compared to T₀F₁ (water foliar application at vegetative growth stage) that was 0.8%.

Table 3: Effect foliar application Zn, Mn and (Zn + Mn) on concentration N in tubers %

Mineral application Time of application	T ₀	T ₁	T ₂	T ₃	Mean
	F ₁	0.80	1.10	1.16	1.48
F ₂	0.85	1.04	1.21	1.39	1.12
F ₃	0.94	1.07	1.13	1.35	1.12
Mean	0.85	1.07	1.16	1.40	
L.S.D 0.05%	T		F	T*F	
	0.04		0.37	0.07	

Phosphorus concentration in tubers:-

Statistical analysis results table 4. showed the significant effect of foliar application of Zn and Mn separately and (Zn + Mn) together on the concentration of p in tubers where T₃ foliar application (Zn + Mn) was superior all other foliar treatment, also it gave the highest concentration of p at 0.28% an increase of 39.81% at an increases of 15.20% when compared to (Zn foliar application) at 0.25% and increase of 2.13% compared to T₂ (Mn foliar application at 0.28% concentration), Time of application has significant effect on p concentration tuber, where f₃ treatment (tuber bulking stage) was superior on all time application, also it gave the highest concentration of p 0.27% in an increase of 14.89% as compared to F₁ foliar application at shoot stage, F₂ foliar application at tuber initiation stage was superior on F₁ in p concentration at 0.27% and increase of 17.78%. Interaction between (Zn + Mn) application and time of application has a significant impact increasing p concentration in tuber, where T₃F₁(application of Zn + Mn at shoot growth stage) was superior on all other interaction treatments and gave the highest p concentration in tuber at 0.33%.

Table 4: Effect foliar application Zn, Mn and (Zn + Mn) on concentration P in tubers %

Mineral application Time of application	T ₀	T ₁	T ₂	T ₃	Mean
	F ₁	0.170	0.260	0.280	0.330
F ₂	0.210	0.240	0.290	0.320	0.265
F ₃	0.238	0.250	0.277	0.315	0.270
Mean	0.206	0.250	0.282	0.288	
L.S.D 0.05%	T		F	T*F	
	0.006		0.005	0.010	

Potassium concentration in tubers

Table 5 shows the significant effect of foliar fertilization of Zn + Mn each separately and together with sprinkling periods and their interaction on the concentration of potassium in tubers of potatoes where T₃ treatment (60 mg Zn L⁻¹ + 30 mg Mn L⁻¹ sprinkling) has given higher potassium concentration in tubers at 2.24% with an increase of 27.27 % compared to least concentration in control treatment F₀ at 1.76 %, Also Mn foliar treatment T₂ was superior as compared to T₁ Zn foliar treatment gave higher potassium in at 2.05 % and 6.77 % increase percentage as compared to potassium concentration in tubers of Zn foliar application at 1.92%, data of foliar application effected was significant in potassium concentration in tubers where foliar application

at vegetative growth was superior potassium concentration at 2.03%, 2.53% and 3.57% increase when compared to potassium concentration in tubers at tuber initiation and tuber bulking stages at 1.98 % and 3.57 % potassium concentration respectively, there was no significant difference between both foliar application at tuber initiation and tuber bulking stages in concentration of potassium in tubers, Interaction Zn and Mn with data of foliar application was significant in potassium concentration in tubers where T₃F₁ (Zn + Mn) foliar application at vegetative growth) by giving potassium concentration in tubers at 2.40 % which 41.18 % increase compared to least concentration of potassium in tubers in interaction treatment T₀F₀ at 1.70 %.

Micronutrient elements play a critical role in plant that lead to increase of leaf area index and there by increased light absorption and increase the amount of dry matter accumulation and economic yield (Ravi et al., 2008), Foliar fertilization not only improves plant growth traits, crop yields and nutrient uptake by crops (Maitlo et al., 2006) but also enhances nutrient use efficiency of crops (Fageria et al., 2009), Numerous studies separately have reported that utilization is increasing performance and quality of potato tubers (Ranjbar and Malakoty, 2000; Mohamadi, 2000; Mousavi et al., 2007),

Table 5: Effect foliar application Zn, Mn and (Zn + Mn) on concentration K in tubers %

Mineral application Time of application	T ₀	T ₁	T ₂	T ₃	Mean
F ₁	1.70	1.98	2.07	2.40	2.03
F ₂	1.76	1.89	2.10	2.18	1.98
F ₃	1.82	1.90	2.00	2.14	1.96
Mean	1.76	1.92	2.05	2.24	
L.S.D 0.05%	T		F	T*F	
	0.04		0.04	0.08	

Kelling and Speth (2001) reported that utilization of elements like Zn and Mn together form resource sulfate Zn and Mn increased and quality of potato crop. Mohamadi (2000) found that application of Zn along with Mn to from foliar application caused increase in efficiency and quality of potato crop. Several researched indicated a positive influence of micronutrient (Zn, Mn) application increase of yield and quantitative parameters of crops (Mosavi et al., 2007) on potato. Al-Fadlly (2016) Found the mean weight of tuber, mean yield tuber per plant and total yield tuber increased when spread with Zn and Mn at vegetative stage.

CONCLUSION

Foliar application of Zn and Mn at (60 ppm Zn + 30 ppm Mn) gave the highest rate of N and K concentration in potatoes tuber at vegetative stage and highest rate of K concentration at bulking stage.

REFERENCES

- [1] Abou El-nour, E. A. A., 2002. Can supplemented potassium foliar feeding reduce the recommended soil potassium? Pak. J. Biol. Sci. 5:259-262.
- [2] AL- Fadlly J. T. M., 2011. Effect of Organic and Mineral fertilizer on growth and yield of potato plants. PHD Dissertation, Collage of Agric, Univ. of Baghdad.
- [3] AL- Fadlly, J. T. M. 2016. Response of potato (*Solanum Tuberosum*) to Foliar Application of Zinc And Manganese Which Fertilized by Organic Fertilized, Journal of Agriculture and Veterinary Science. 9(4):87-91
- [4] Bremner, J. M., and Mulvaney, C. S. 1982. Nitrogen – Total. In: Methods of Soil Analysis, Part 11. Chemical and Microbiological Properties, eds, A. L. Page, R. H. Miller, and D. R. Keeney, pp. 595 – 624. Madison, W1: American Society of Agronomy.
- [5] Clarkson, D. T., 1988. The uptake and translocation of manganese by plant roots. In: R.D. Graham; R. J. Hannam; N. J. Uren (eds). Manganese in Soil and Plants. Kluwer. Acadmic Publishers, Dordrecht, The Netherlands, pp: 101-111.

- [6] Eppendorf, N. and G. Hing. 1970. Interaction manual of flame photometer B 700-E. Measuring method, Description of apparatus and Instructions for use.
- [7] Fageria, N. K., Filho, M. P. B., Moreira, A., and C. M. 2009. Foliar fertilization of crop plants. *Journal of Plant Nutrition*, 32 (6), 1044 – 1064.
- [8] Flis B., Zimnoch-Guzowska E., Mankowski D., 2012. Correlation among yield, taste, tuber characteristics and mineral contents of potato cultivars growing conditions. *Journal of Agricultural Science*. 4(7), 197-207.
- [9] Grotz, N, and Guerinot, M. L. 2006. Molecular aspects of Cu, Fe and Zn homeostasis in plants. *Biochimica et Biophysica Acta*. 1763(7):595-608.
- [10] Jackson, C., J. Dench, A. L. Moore, B. Haliwell, C. H, Foyer and D. O. Hall, 1978. Sub cellular location and identification of super oxide dismutase in leaves of higher plants. *European Journal of Biochemistry*, 91:339-344.
- [11] Kaya, C. and D. Higgs, 2002. Response of tomato (*Lycopersicon esculentum* L.) cultivars to foliar application of zinc when grown in sand culture at low zinc. *Sci. Hort.*, 93: 53 – 64.
- [12] Kelling, K. A. and P. E. Speth, 2001. Effect of micronutrient on potato tuber and quality at Spooner, 2001. Department of soil Science - University of Wisconsin - Madis.
- [13] Maitlo, A., Zia Ulhassan, A., Shah, N., and Khan, H. 2006. Growth, yield and nutrient uptake of wheat (*Triticum aestivum*) in relation to foliar and soil application of urea. *International Journal of Agriculture And Biology*, 8, 477 – 481.
- [14] Millaleo, R., M. Reyes-Díaz., A. G. Ivanov, M. I. Mora and M. Alberdi, 2010. Manganese as essential and toxic element for plants: transport, accumulation and resistance mechanisms. *J. Soil. Sci. Pant Nutr.*, 10(4): 476-494.
- [15] Mohamadi, E. 2000. Study Effect of nutrient elements utilization methods (Zn, Mn, and Mg) on increase performance quantitative and quality of two potato species. Jihad and Agriculture Ministry. Final Report of Research Institute Reformand Providing Sapliny and seed.
- [16] Mousavi, S. R., Galavi. M., Ahmadvand, G., 2007. Effect of Zinc and Manganese Foliar Application on Yield, Quality and Enrichment on Potato (*Solanum tuberosum* L.) *Asian, J, Plant Sci.*, 6:1256 – 1260.
- [17] Muharem, Hussein jawad and Karim Saleh Abdul, 1987. The effect of planting date and source of tuber on the quality of potato tubers in autumn and spring Aerutin hid in Xbat / Erbil region. 5(4):33-37.
- [18] Mukhopadhyay, M. and A. Sharma, 1991. Manganese in cell metabolism of higher plants. *Bot. Rev.*,57:177-149.
- [19] Ranjbar R, Malakoty, M. J. 2000. Effect of different amount K and sulfate Zn fertilizers on performance and quality of potato (*Solanum tuberosum* L.) crop in Banab (first and second part). *J. Sci. Res. Soil and water, Natl. Soil Water Res. Inst. Tehran, Iran*, 12: 110-120.
- [20] Ravi, S, Channal, HT, Hebsur, NS, Patil, BN, Dhamatti, P. R. 2008. Effect of sulphur, zinc and iron nutrition on growth, yield, nutrient uptake and quality of safflower (*Carthamus tinctorius* L.). *Karnataka, J. Agric. Sci.*, 21(3):382 – 385.
- [21] SAS, 2001. User guide ststistic (Version 6-12). SAS Inst. Cary, N. C. USD.
- [22] Uygur, V. and D. L. Rimmer, 2000, Reaction of zinc with iron coated calcite surface at alkaline pH. *Eur. J. Soil. Sci.*, 51:511-516.
- [23] Watanable, F.C. and S. R. Olsen, 1965. Test of ascorbic acid method for determining phosphorus in water and NaHCO₃ extracts from soils, *Soil Sci. Soc. Am. Proc.*, 29:677-6