



Research Journal of Pharmaceutical, Biological and Chemical Sciences

Kinetics and Equilibrium Studies on Removal of Cd²⁺ From Aqueous Solution on to *Bacillus Subtilis*

Sethuraman Pandiyan and Dharmendirakumar Mahendradas

Department of Chemical Engineering, A.C.Tech Campus, Anna University, Chennai-600 025, India

ABSTRACT

Biosorption of heavy metals can be an effective process for the removal of toxic heavy metals present in the wastewater. An attempt is made in the present investigation to remove Cd²⁺ ions from aqueous solutions using *B.subtilis*. Batch experiments were carried out for Cd²⁺ removal over a wide range of operating conditions. It has been noticed that the Cd²⁺ removal capacity dependent on the initial pH and the initial Cd²⁺ concentration. Experimental data were analyzed with kinetics and isotherm models. It has been observed that the pseudo second order kinetics and Freundlich isotherm equilibrium model fit well with the present data. The FTIR analysis of bacterial biomass revealed the presence of amino, carboxyl, hydroxyl and carbonyl groups, which are responsible for biosorption of Cd²⁺ metal ions. The results indicated that the biomass of *B.subtilis* is an efficient bioaccumulant for the removal of Cd²⁺ ions from aqueous solutions.

Keywords: Biosorption; Cd²⁺ removal; Kinetics; Isotherms; *Bacillus subtilis*-live and dead

***Corresponding author**

Email: mdkumar@annauniv.edu



INTRODUCTION

The growth of industries and day to day changes in human activities has resulted in the increase in volume and complexity of wastewater in the environment. Industries such as metal plating & tanneries uses heavy metal in their processes which generates large amount of aqueous effluents with high levels of heavy metals. The heavy metals include cadmium, chromium, cobalt, copper, iron, manganese, mercury, molybdenum, nickel, silver, and zinc. Heavy metal pollution poses serious environmental issue due to its detrimental effect on human health. The removal and recovery of heavy metals from wastewater becomes important to protect the environment and human health. Cd^{2+} is one such heavy metal responsible for polluting the ecosystem. It is used in a wide variety of industrial processes such as alloy preparation, metal plating and electronics. Cd^{2+} effectively binds in human body by high molecular protein such as albumin and nonprotein sulfhydryl group. This is accumulated in the kidneys and liver. Excess Cd^{2+} in the organisms can damage DNA sequencing and may cause genetic changes and cancer [1]. It also appears to be the largest single contributor to autoimmune thyroid disease [2]. Unlike organic pollutants, which can be degraded, metallic pollutants are immutable and will present indefinitely in the environment [3]. The non biodegradable nature and long biological half-life of most metals lead to potential accumulation and human exposure via water or food [4]. Lead and zinc mines are the main sources for the release of lead, chromium and cadmium into the natural environment [5].

The traditional methods used for the removal of Cd^{2+} present in the effluent include precipitation, evaporation, adsorption on activated carbon, ion exchange membrane processing, and solvent extraction. However, these methods are expensive and inefficient to remove metals at low concentration [6-8]. The use of biological material, including living and non-living microorganisms, in the removal and possible recovery of toxic or precious metals from industrial wastes, has gained greater important in the recent years. The major advantages of the biosorption technology are its effectiveness in quickly reducing the concentration of heavy metal ions to very low levels with high efficiency by using inexpensive biosorbent materials [9].

The physicochemical interactions between metal ions and different functional groups on the biomass surface such as carboxyl, hydroxyl, sulfhydryl and amino groups play an important role in the biosorption process. Living, dead and immobilized cells can be utilized in this process. Moreover, the regeneration of biosorbent for multiple uses is easy as; it shows selectivity towards different metals removal. Bacteria have a high surface area-to-volume ratio that can provide a large contact interface, and this allows the interaction with metals from the surrounding environment. The advantages of biological substrates include diversity of biologically active binding sites and less interference from alkali/alkali-earth metals [10]. Microorganisms comprises of bacteria, algae, fungi and yeast which uptake metals either actively (bioaccumulation) and/or passively (biosorption) [11, 12]. More recently, such as bacteria microbial biomass is specifically considered useful for removal of heavy metals from aqueous solutions [13, 14].

In this study, we evaluate the effectiveness in the removal of Cd^{2+} ions from aqueous solution through biosorption by dead and live *B.subtilis* and also to characterize the metal biosorption behavior of hyper resistant *B.subtilis* strain for Cd^{2+} ion. Pseudo first & second order kinetics and Langmuir & Freundlich isotherm parameters were also evaluated from batch biosorption assay. Bacteria *B.subtilis* is an alternative low-cost bioadsorbent for successful removal of Cd^{2+} from aqueous solutions. The high amount of Cd^{2+} uptake by *B.subtilis* places this biosorbent as the best adsorbent for this metal removal from aqueous solution.

MATERIALS AND METHODS

Preparation of the powdered dried dead cells

B.subtilis (MTCC-121) species were purchased from Microbial type culture collection, Chandigarh, India. Nutrient broth culture media was prepared and maintained as per the guidelines of MTCC. The slant cultures were prepared with prescribed growth medium containing beef extract 1.0g, yeast extract 2.0g, peptone 5.0g, sodium chloride 5.0g and distilled water 1.0 litre. The culture was sterilized in an autoclave maintained at 15 lbs for 15 minutes. The chemical Cadmium Sulphate ($3\text{CdSO}_4 \cdot 8\text{H}_2\text{O}$) used in the present investigation was of analytical grade purchased from Ranbaxy Fine Chemicals Ltd., India. Stock solution of Cd^{2+} (1000 mg/l) was prepared using purified double distilled water. Cd^{2+} solutions of varying concentration were obtained by diluting the stock solution. The Cd^{2+} concentration was analyzed using an atomic absorption spectrophotometer (AAAnalyst 800; PerkinElmer, USA). The morphology of bacteria has been analyzed using Scanning electron microscopy (JSM-6360; JEOL, Japan). Fourier transform infrared spectroscopy (FTIR; Model Tensor 27, Bruker Optic GmbH, Germany) spectrometer was used to determine the type of functional groups in bacteria responsible for the Cd^{2+} metal accumulation.

Biosorption of Cd^{2+} ion

The biosorption studies have been carried out with monometallic solutions prepared from stock solutions in a batch system. A known quantity of metal ion concentration solution was taken in flasks and biomass was added. The flasks were shaken at 150 rpm in 30 °C for a certain time of 0–24 hrs. Test samples were collected at regular intervals of time and centrifuged at 6000 rpm for 10 min, filtered through 0.2 μm filter and analyzed for Cd^{2+} concentration using AAS.


RESULTS AND DISCUSSION

Effect of Growth curve

During lag phase, *B.subtilis* adapt themselves to grow in a particular condition with closed system. This is the period where the individual *B.subtilis* are maturing and not yet able to

divide. During the lag phase of the bacterial growth cycle, synthesis of RNA, enzymes and other molecules occurs. Exponential phase or log phase is a period characterized by cell doubling. The number of new bacteria appearing per unit time is proportional to the present population. If growth is not limited, doubling will continue at a constant rate so both the number of cells and the rate of population doubles with each consecutive time period. During stationary phase, the growth rate slows as a result of nutrient depletion and accumulation of toxic products. This phase is reached as the bacteria begins to exhaust the resources that are available to them. This phase has a constant rate value as the rate of bacterial growth is equal to the rate of bacterial death. At death phase, bacteria runs out of nutrients, no availability of space and Oxygen lead to death. The growth rate of *B.subtilis* has been recorded for lag phase to dead phase and the observation is given in Figure 1. It can be ascertained that the lag phase exists for first 4 hrs followed by a consistent exponential pattern of increase in log phase for 12 hrs. The stationary phase exists for 20 hrs.

Figure 1: Growth curve of live *B.subtilis* in closed system optical density at 600 nm.



Isotherms

In biosorption, the metal ions accumulate and the surface of bacterium cell wall which can be well represented by a conventional accumulation isotherm. An attempt is made to test the Langmuir and Freundlich isotherms models for metal removal.

Langmuir isotherm

The Langmuir isotherm assumes monolayer adsorption on homogenous biosorbent surface (cell wall). The mathematical expression of Langmuir isotherm can be given as

$$q_e = \frac{K_L b C_e}{1 + b C_e} \quad (3)$$

The linearization of the above equation results in:


$$\frac{C_e}{q_e} = \frac{1}{K_L} + \frac{a_L}{K_L} C_e \quad (4)$$

The binding constant K_L , and the sorbent capacity a_L are estimated by plotting C_e/q_e against C_e . The model simulations along with experimental observations for Cd^{2+} with the experimental values of K_L and a_L along with the linear regression co-efficient (R^2) are given Table 1 and Figure 2 respectively.

Table 1: Assignments of Infrared absorption bands

Wave numbers (cm-1)	Intensity shape	Assignment
3600-3750	Sharp	O-H stretching
3400-3550	Sharp	O-H stretching
3100-3500	Strong-broad	N-H stretching
2500-3400	Weak-broad	O-H stretching
2700-2950	Variable	C-H stretching
1400-1660	Variable	N-H bending
1280-1430	Variable	C-H bending
1160-1420	Variable	O-H bending
900-1350	Variable	C-N stretching
900-1380	Variable	C-O stretching
800-880	Medium-strong	N-H and C-H rocking

Figure 2: Langmuir isotherms of Cd^{2+} ion on live and dead biomass of *Bacillus subtilis*.



Freundlich isotherm

The Freundlich isotherm is an empirical model that relates the accumulation intensity of the sorbent to accumulants. The isotherm is adopted to describe reversible accumulation and

not restricted to monolayer formation. The mathematical expression of the Freundlich isotherm model can be given as

$$q_e = K_F C_e^{b_F} \quad (5)$$

Where K_F and b_F are the constants which give accumulation capacity and accumulation intensity respectively. A linear form of the Freundlich model can be written as follows


$$\log q_e = \log K_F + b_F \log C_e \quad (6)$$

A plot of $\log q_e$ versus $\log C_e$ gives a straight line with slope K_F and intercept b_F . The values of K_F and b_F along with the linear regression co-efficient (R^2) for the present experimental conditions have been obtained and are given in Table 2 and Figure 3. It can be observed from the correlation coefficient (R^2) that the Freundlich isotherm model matches satisfactorily with the experimental observation.

Table 2: Langmuir and Freundlich isotherms constants for the biosorption of Cd^{2+} on live and dead of *Bacillus subtilis*.

Metal	Langmuir constant			Freundlich constant		
	K_L	b	R^2	K_F	b_F	R^2
Live	0.003	2.861	0.889	1.563	8.796	0.993
Dead	0.011	2.401	0.998	1.216	3.505	0.998

Figure 3: Freundlich isotherm of Cd^{2+} ion on live and dead biomass of *Bacillus subtilis*.



Adsorption Kinetics models

In the sorption studies, equilibrium of sorption and sorption kinetics are two important physicochemical factors to be considered. Sorption kinetics can explain the dependency of

sorption rates upon the concentrations of biosorbate in solution, and how sorption rates are affected by sorption capacity, or by the character of the sorbent [16]. Since the principles underlying biosorption kinetics include fitting the model that represents the experimental data best [17], the pseudo-first-order and the pseudo second order kinetic models were tested in our study.

Pseudo first order model

Pseudo first order kinetic model by the following equation

$$\log(q_e - q_t) = \log(q_e) - \frac{k_1 t}{2.303} \quad (3)$$

In place where q_e and q_t are the Cd^{2+} ions where sorbed at equilibrium and time t , respectively. K_1 is the rate constant for pseudo first order biosorption. This model was successfully applied to describe the kinetics for many adsorption systems. The kinetic data in Table 3 demonstrate that the biosorption of Cd^{2+} onto mixed biosorbent follow the negative pseudo first order kinetics.

Table 3: Kinetics data for the uptake by live and dead of Bacillus subtilis.

Metal	Pseudo first order			Pseudo second order		
	q_e	K_1	R^2	q_e	K_2	R^2
Live	0.194	0.009	0.209	0.030	50.00	0.972
Dead	0.150	0.1087	0.289	0.013	370.37	0.996


Pseudo second order model

The adsorption kinetics in addition described by pseudo second order model. The linearized integral form of the model by the following equation

$$\frac{t}{q} = \frac{1}{k_2 q_e^2} + \frac{t}{q_e} \quad (4)$$

In Position k_2 is the pseudo second order rate constant and q_2 is the equilibrium biosorption capacity. Values of k_2 and q_2 were calculated from a plot of t/q_t against t Figure 4. From Table 3 the (R^2) values for the pseudo first order and the pseudo second order kinetic models indicated that the pseudo first order kinetic model could not explain the biosorption kinetics in our study. The values of equilibrium biosorption capacities from the pseudo second order kinetic model with (R^2) values 0.972 & 0.996 for live & dead respectively.


Figure 4: Pseudo second order kinetic modeling of Cd²⁺ adsorption on biomass of Bacillus subtilis.



Fourier transform infrared spectroscopy

The FTIR spectra of B.subtilis biomass before and after the metal uptake are shown in Figure 5.

Figure 5: Infra spectra of (a) live B.subtilis were treated with Cd²⁺ ions for different lengths of time namely (b) 0.0, (c) 0.2, (d) 0.12, (e) 0.24 hrs, (f) dead B.subtilis and (g) dead Cd²⁺ ion.





It can be ascertained that the figure indicates the presence of amino, carboxylic, hydroxyl and carbonyl groups. It can be noticed from the figure that a strong broad and intense peak at $3400\text{-}3700\text{ cm}^{-1}$ shows the presence of hydrogen-bonded hydroxyl group. This also coupled with stretching vibration of the NH_2 moiety (Table 1). A strong absorption at 1050 cm^{-1} shows the stretching of C-O in polysaccharides [15]. Both live and dead *B.subtilis* were subjected for Cd^{2+} accumulation. Samples were taken at regular intervals and analyzed by using FTIR spectra. A peak at 1280 and 1470 cm^{-1} has been observed for all intervals of time which shows the presence of carboxylic and amid groups. While the peak at 885 and 618 cm^{-1} during the first 12 hrs represent the stretching and bending modes. It can be noticed from figure that the IR spectrum for all intervals of time except 24 hrs shows a peak with strong vibration at 1280 and 1470 cm^{-1} , represents presence of C-H bonding. Finally the peaks at $2950\text{-}3000\text{ cm}^{-1}$ with CH_2 stretching frequency shows the participation of OH and NH_2 . The analysis of the FTIR spectra showed the presence of ionisable functional groups (carboxyl, amino, amide and hydroxyl) able to interact with protons or metal ions. The above results obtained give an idea about the presence of functional groups on the bacterial cell surfaces.

Scanning electron microscopy

The Scanning electron microscopy (SEM) images clearly reveal that the surface texture and morphology of the biosorbent are shown in Figure 6 and 7, which represent with and without Cd^{2+} ions, with different magnifications. The length and width of the filament of live *B.subtilis* varies from 1.0 to $2.5\text{ }\mu\text{m}$ and 0.3 to $0.8\text{ }\mu\text{m}$ and as expected the cell shrinks when it is dried. In dry species the length and width of the bacilli lies in the range of 0.6 to $1.8\text{ }\mu\text{m}$ and 0.1 to $0.4\text{ }\mu\text{m}$. The difference in the surface morphology after the metal uptake by live and dead species is evident from SEM images. The surface of the biomass becomes rough after metal uptakes. It has already been stated in the preceding section on IR spectroscopy and kinetic study that live *B.subtilis* takes up more metal ions than the dead one. A close examination of the SEM micrograph of the treated samples also supports this fact; the width of the cell of treated live biomass is more compared to the same for dried species. When treated with metal of Cd^{2+} ions there is the possibility of transport of Cd^{2+} ions through the cell membrane in live biomass that may be responsible for more uptakes in this *Bacillus subtilis*. The SEM micrograph of metal-treated live *B.subtilis* shows metal ions protruding in the strand of the cell, supporting membrane transport process.

Figure 6: SEM Images represent live bacilli A and dead bacilli B before accumulation on *Bacilli subtilis* without the presence of Cd^{2+} .



CONCLUSION

The presented study is based on the biosorption of *B.subtilis* on Cd^{2+} ions by live and dead biomass. The metabolic activities in live *B.subtilis* possibly helps in higher uptake of metal ions compared to dead species. The sorption process for live biomass using Freundlich model given a good correlation coefficient and in the case of dead biomass both isotherms are obeyed. The uptake of Cd^{2+} ions by live and dead *B.subtilis* follows pseudo second-order rate equation. The Infra red spectra suggest the participation of carboxylate, hydroxyl of polysaccharides, phosphate and amide groups in the uptake. SEM and microscopic studies support the sorption data. Consequently, removal of Cd^{2+} live and dead *B.subtilis* took up significant amount of Cd^{2+} ions. In this live *B.subtilis* were found to be simple, fast and efficient.

REFERENCES

- [1] Carmichael W. The toxins of cyanobacteria. *Scientific American* 1994; 170:78–86.
- [2] Der R, Yousef M, Fahim Z, Fahim M. *Chem Pathology and Pharm* 1977; 17:237–253.
- [3] Sharma P, Kumari P, Srivastava MM, Srivastava S. *Biores Tech* 2006; 97:299–305.
- [4] Isabel V, Nuria F, Maria M, Nuria M, Jordi P, Joan S. *Water Res* 2004; 38:992–1002.
- [5] Liu H, Probst A, Liao B. *Science of the Total Envi* 2005; 339:153–166.
- [6] Rickelton WA. *Hydrometallurgy* 1998; 50:339–344.
- [7] Wombacher F, Rehkammer M, Mezger K. *Geochim Cosm Acta* 2004; 68:2349–2357.
- [8] Selatnia A, Bakhti MZ, Madani A, Kertous L, Mansouri Y. *Hydrometallurgy* 2004; 75:11–24.
- [9] Volesky B. *Sorption and Biosorption*. BV Sorbex, Inc, Montreal, Canada 2003; 316.
- [10] Madrid Y, Camara C. *Trends in anal chem* 1997; 16:36–44.
- [11] Hussein H, Ibrahim SF, Kandeel K. *E Journal of Biotech* 2004; 1:38–46.
- [12] Yan G, Virarghavan T. *Water South Africa* 2000; 26:119–123.
- [13] Nourbakhsh M, Kilic S, Arslan Ilhan S, Ozdag H. *Chem Eng J* 2002; 85:351– 55.
- [14] Ahluwalai SS, Goyal D. *Biores Tech* 2007; 98:2243–2257.
- [15] Rathinam A, Balaraman M, Rao JR, Nair BU, Thirumachari R. *Envir Sci & Tech* 2004; 38:300–306.
- [16] Ho YS, Ng JCY, McKay G. *Separ and Purifi Meth* 2000; 29:189–232.
- [17] Kuppasamy V, Joseph RJ, Kandasamy P, Manicakam V. *E Journal of Biotech* 2004; 7:61–71.