

Research Journal of Pharmaceutical, Biological and Chemical Sciences

Impact of Distillery Spentwash Irrigation on Sprouting and Growth of *Anthurium* (Araceae) Flowering plant

Chandraju S^{1*}, Thejovathi C¹, and Chidan Kumar CS²

¹ Departmentt of Studies in Sugar Technology, Sir M. Vishweswaraya Postgraduate Center, University of Mysore, Tubinakere, Mandya -571402, Karnataka, India

² Department of Chemistry, Alva's Institute of Engineering & Technology, Mijar, Moodbidri-574225, South Canara Dt. Karnataka, India

ABSTRACT

Sprouting and growth of *Anthurium* (Araceae) flowering plant was made by irrigated with distillery spentwash of different concentrations. The spentwash i.e., primary treated spentwash (PTSW), 1:1, 1:2, and 1:3 spentwash were analyzed for their plant nutrients such as nitrogen, phosphorous, potassium and other physical and chemical characteristics. Experimental soil was tested for its chemical and physical parameters. *Anthurium* (Araceae) sets were planted in different pots and irrigated with raw water (RW), 1:1, 1:2 and 1:3 spentwash. The nature of sprouting and growth was studied. It was found that the sprouting and growth of plant was very good (100%) in 1:3 SW irrigation, while very poor (25%) in 1:1 SW, moderate (80%) in 1:2 SW and 95% in RW irrigation growth. This concludes that the diluted spentwash can be conveniently used for irrigation purpose without adverse affect on soil.

Keywords: *Anthurium*, irrigation, spentwash, Araceae

*Corresponding author

INTRODUCTION

Anthurium (Araceae) is a large genus of about 600-800 species [1], belonging to the arum family (Araceae). *Anthurium* can also be called “Flamingo Flower” or “Boy Flower”, both referring to the structure of the spathe and spadix. Most species occur in Panama, Colombia, Brazil, the Guiana Shield and Ecuador. According to the work of noted aroid botanist Dr. Tom Croat of the Missouri Botanical Garden, no members of this genus are indigenous to Asia [2]. Deliberately or accidentally, however, some species have been introduced into Asian rain forests, and have become established there as aliens. *Anthurium* grows in many forms, mostly evergreen, bushy or climbing epiphytes with roots that can hang from the canopy all the way to the floor of the rain forest. There are also many terrestrial forms which are found as understory plants, as well as hemi epiphytic forms. The stems are short to elongate with a length between 15 and 30 cm. The simple leaves come in many shapes; most leaves are to be found at the end of the stems, although terrestrial plants show less of this pachycaul tendency. Leaves may be spatulate, rounded, or obtuse at the apex. The leaves are petiolate and possess a structure called the reticulum, which is unique to the genus *Anthurium*. *Anthurium* flowers are small and develop crowded in a spike on a fleshy axis, called a spadix, characteristics of the Araceae. The flowers on the spadix are often divided with a sterile band separating male from female flowers. This spadix can take on many forms and colors. The flowers of *Anthurium* give off a variety of fragrances, each attracting a variety of specific pollinators. Several species are popular in the florist trade as pot plants or cut flowers and for interior decoration.

Molasses (one of the important byproducts of sugar industry) is the chief source for the production of ethanol in distilleries by fermentation method. About 08 (eight) liters of wastewater is generated for every liter of ethanol production in distilleries, known as raw spent wash (RSW), which is known for high biological oxygen demand (BOD: 5000-8000mg/L) and chemical oxygen demand (COD: 25000-30000mg/L), undesirable color and foul odor [3]. Discharge of RSW into open field or nearby water bodies results in environmental, water and soil pollution including threat to plant and animal lives. The RSW is highly acidic and contains easily oxidisable organic matter with very high BOD and COD [4]. Also, spentwash contains high organic nitrogen and nutrients [5]. By installing biomethanation plant in distilleries, reduces the oxygen demand of RSW, the resulting spentwash is called primary treated spentwash (PTSW) and primary treatment to RSW increases the nitrogen (N), potassium (K), and phosphorous (P) contents and decreases calcium (Ca), magnesium (Mg), sodium (Na), chloride (Cl⁻), and sulphate (SO₄²⁻) [6]. PTSW is rich in potassium (K), sulphur (S), nitrogen (N), phosphorous (P) as well as easily biodegradable organic matter and its application to soil has been reported to increase yield of sugar cane, wheat and rice [7], Quality of groundnut [8] and physiological response of soybean [9]. Diluted spentwash could be used for irrigation purpose without adversely affecting soil fertility [10], seed germination and crop productivity [11]. The diluted spentwash irrigation improved the physical and chemical properties [12] of the soil [13] and further increased soil micro flora [14]. Twelve pre-sowing irrigations with the diluted spentwash had no adverse effect on the germination of maize but improved the growth [15]. Diluted spent wash increases the growth of shoot length, leaf number per plant, leaf area and chlorophyll

content of peas [16]. Increased concentration of spentwash causes decreased seed germination, seedling growth and chlorophyll content in Sunflowers (*Helianthus annuus*) and the spentwash could safely used for irrigation purpose at lower concentration [17]. The spent wash contained an excess of various forms of cations and anions, which are injurious to plant growth and these constituents should be reduced to beneficial level by diluting spentwash, which can be used as a substitute for chemical fertilizer [18]. The spentwash could be used as a complement to mineral fertilizer to sugarcane [19]. The spentwash contained N, P, K, Ca, Mg and S and thus valued as a fertilizer when applied to soil through irrigation with water [20]. The application of diluted spentwash increased the uptake of Zinc (Zn), Copper (Cu), Iron (Fe) and Manganese (Mn) in maize and wheat as compared to control and the highest total uptake of these were found at lower dilution levels than at higher dilution levels . Mineralization of organic material as well as nutrients present in the spentwash was responsible for increased availability of plant nutrients. Diluted spentwash increase the uptake of nutrients, height, growth and yield of leaves vegetables [21], nutrients of cabbage and mint leaf [22], nutrients of top vegetable [23], pulses, condiments, root vegetables, of some root vegetables in untreated and spentwash treated soil, yields of top vegetables (creepers). However, no information is available on sprouting and growth of *Anthurium* flowering plant irrigated by distillery spentwash. Therefore, the present investigation was carried out to study the influence of different proportions of spentwash on the sprouting and growth of *Anthurium*.

MATERIALS AND METHODS

Physio-chemical parameters [24] and amount of nitrogen (N) [25], potassium (K), [26] phosphorous (P) [27] and sulphur (S) [28] present in the primary treated diluted spentwash (1:1, 1:2 and 1:3 SW) were analyzed by standard methods [29]. The PTSW was used for irrigation with a dilution of 1:1, 1:2 and 1:3. A composite soil sample collected prior to spentwash irrigation was air-dried, powdered and analyzed for physico-chemical properties [30]. Flowering [31] plants [32] selected for the present investigation were *Anthurium*. The sets were planted in different pots (30(h), 25(dia)) and irrigated (by applying 5-10mm/cm² depends upon the climatic condition) [33] with raw water (RW), 1:1 SW, 1:2 SW and 1:3 SW at the dosage of twice a week and rest of the period with raw water as required. Cultivation was conducted in triplicate, in each case sprouting, growth were recorded.

Table 1: Chemical characteristics of distillery Spentwash

Chemical parameters	PTSW	1:1 PTSW	1:2 PTSW	1:3 PTSW
pH	7.57	7.63	7.65	7.66
Electrical conductivity ^a	26400	17260	7620	5330
Total solids ^b	47200	27230	21930	15625
Total dissolved solids ^b	37100	18000	12080	64520
Total suspended solids ^b	10240	5380	4080	1250
Settleable solids ^b	9880	4150	2820	3240
COD ^b	41250	19036	10948	2140
BOD ^b	16100	7718	4700	2430
Carbonate ^b	Nil	Nil	Nil	Nil
Bicarbonate ^b	12200	6500	3300	1250

Total Phosphorous ^b	40.5	22.44	17.03	10.80
Total Potassium ^b	7500	4000	2700	1620
Calcium ^b	900	590	370	190
Magnesium ^b	1244.16	476.16	134.22	85
Sulphur ^b	70	30.2	17.8	8.4
Sodium ^b	520	300	280	140
Chlorides ^b	6204	3512	3404	2960
Iron ^b	7.5	4.7	3.5	2.1
Manganese ^b	980	495	288	160
Zinc ^b	1.5	0.94	0.63	0.56
Copper ^b	0.25	0.108	0.048	0.026
Cadmium ^b	0.005	0.003	0.002	0.001
Lead ^b	0.16	0.09	0.06	0.003
Chromium ^b	0.05	0.026	0.012	0.008
Nickel ^b	0.09	0.045	0.025	0.012
Ammonical Nitrogen ^b	750.8	352.36	283.76	178
Carbohydrates ^c	22.80	11.56	8.12	6.20

Units: a – μ S, b – mg/L, c- %, PTSW - Primary treated distillery spentwash

Table 2: Amount of N, P, K and S (Nutrients) in distillery Spentwash

Chemical parameters	PTSW	1:1 PTSW	1:2 PT SW	1:3 PTSW
Ammonical Nitrogen ^b	750.8	352.36	283.76	160.5
Total Phosphorous ^b	40.5	22.44	17.03	11.2
Total Potassium ^b	7500	4000	2700	1800
Sulphur ^b	70	30.2	17.8	8.6

Unit: b – mg/L, PTSW - Primary treated distillery spentwash

Table 3: Characteristics of experimental soil

Parameters	Values
Coarse sand ^c	9.24
Fine sand ^c	40.14
Slit ^c	25.64
Clay ^c	20.60
pH (1:2 soln)	8.12
Electrical conductivity ^a	530
Organic carbon ^c	1.64
Available Nitrogen ^b	412
Available Phosphorous ^b	210
Available Potassium ^b	110
Exchangeable Calcium ^b	180
Exchangeable Magnesium ^b	272
Exchangeable Sodium ^b	113
Available Sulphur ^b	330
DTPA Iron ^b	204
DTPA Manganese ^b	206
DTPA Copper ^b	10
DTPA Zinc ^b	55

Units: a – μ S, b – mg/L, c- %

Table 4: Characteristics of experimental soil (After harvest)

Parameters	Values
Coarse sand ^c	9.69
Fine sand ^c	41.13
Slit ^c	25.95
Clay ^c	24.26
pH (1:2 soln)	8.27
Electrical conductivity ^a	544
Organic carbon ^c	1.98
Available Nitrogen ^b	434
Available Phosphorous ^b	218
Available Potassium ^b	125
Exchangeable Calcium ^b	185
Exchangeable Magnesium ^b	276
Exchangeable Sodium ^b	115
Available Sulphur ^b	337
DTPA Iron ^b	212
DTPA Manganese ^b	210
DTPA Copper ^b	12
DTPA Zinc ^b	60

Units: a – μ S, b – mg/L,

Table 5: Growth of *Anthurium* plant at different irrigations (cm)

RW 15 th 22 nd 29 th (Day)	1:1SW 15 th 22 nd 29 th (Day)	1:2 SW 15 th 22 nd 29 th (Day)	1:3 SW 15 th 22 nd 29 th (Day)
20, 24, 26	03, 05, 08	26, 28, 30	28, 30, 33

RESULTS AND DISCUSSION

Chemical composition of P₂SW, 1:1, 1:2, and 1:3 SW such as pH, electrical conductivity, total solids (TS), total dissolved solids (TDS), total suspended solids (TSS), settleable solids (SS), chemical oxygen demand (COD), biological oxygen demand (BOD), carbonates, bicarbonates, total phosphorous (P), total potassium (K), ammonical nitrogen (N), calcium (Ca), magnesium (Mg), sulphur (S), sodium (Na), chlorides (Cl), iron (Fe), manganese (Mn), zinc (Zn), copper (Cu), cadmium (Cd), lead (Pb), chromium (Cr) and nickel (Ni) were analyzed and tabulated (Table-1). Amount of N, P, K and S contents are presented (Table-2). Characteristics of experimental soils such as pH, electrical conductivity, the amount of organic carbon, available nitrogen (N), phosphorous (P), potassium (K), sulphur (S), exchangeable calcium (Ca), magnesium (Mg), sodium (Na), DTPA iron (Fe), manganese (Mn), copper (Cu) and zinc (Zn) were analyzed and tabulated (Table-3 & 4). It was found that the soil composition is fit for the cultivation of plants, because it fulfils all the requirements for the growth of plants. Sprouting and growth of *Anthurium* plant leaves, uptakes of all the parameters were very good in both 1:2 and 1:3 spent wash as compared to 1:1, SW and raw water. In both 1:1, 1:2 and 1:3 spent wash irrigation [31], the uptake of the nutrients [32] such as fat, calcium, zinc, copper and vitamins carotene and vitamin c were almost similar but the uptake of the nutrients [33] and parameters such as protein, fiber, carbohydrate, energy, magnesium and phosphorous were much more in the case



of 1:1, 1:2, spent wash irrigation than 1:3, and raw water irrigations (Table-5). This could be due to the more absorption of plant nutrients present in spent wash by plants at higher dilutions. It was also found that no negative impact of heavy metals like lead, cadmium and nickel on the leaves of *Anthurium* plant. The soil was tested after the harvest; found that there was no adverse effect on soil characteristics (Table-4).

CONCLUSION

It is found that the nutrients uptake in the Sprouting and growth of *Anthurium* (Araceae) plant were largely influenced in case of 1:1, 1:2 and 1:3 SW irrigation than with raw water. But 1:3 distillery spentwash shows more uptakes of nutrients when compared to 1:2 SW. This could be due to the maximum absorption of nutrients by plants at more diluted spentwash. After harvest, soil has tested; found that there was no adverse effect on characteristics. Hence the spentwash can be conveniently used for irrigation purpose with required dilution without affecting environment and soil.

ACKNOWLEDGEMENTS

Authors are grateful to The General Manager, Nijaveedu Sugars Limited. Koppa, Maddur Tq., Karnataka, for providing spentwash

REFERENCES

- [1] Sunset Western Garden Book 1995:606-607.
- [2] Abstract (<http://www.aroid.org/genera/Anthurium/abstrap 1.htm>)
- [3] Joshi HC, Kalra N, Chaudhary A, Deb DL. Asia Pac J Environ Develop 1994; 1: 92-103.
- [4] Patil JD, Arabatti SV, Hapse DG. Bartiya sugar 1987; 9-15.
- [5] Ramadurai R, Gerard EJ. Sugar J 1994; 20: 129-131.
- [6] Mohamed Haroon AR, Subash Chandra Bose M. Indian Farm 2004; 48- 51.
- [7] Pathak H, Joshi HC, Chaudhary A, Chaudhary R, Kalra N, Dwivedi MK. J Indian Soc Soil Sci 1998; 46: 155-157.
- [8] Amar BS, Ashisk B, Sivakoti R. J Plant Nutr Soil Sci 2003; 166: 345-347.
- [9] Ramana S, Biswas AK, Kundu S, Saha JK, Yadava RBR. Plant Soil Res 2000; 2: 1-6.
- [10] Kaushik K, Nisha R, Jagjeeta K, Kaushik CP. Bioresource Technol 2005; 96(17): 1860-1866.
- [11] Kuntal MH, Ashis K, Biswas AK, Misra K. Nutr Soil Sci 2004; 167(5): 584-590.
- [12] Raverkar KP, Ramana S, Singh AB, Biswas AK, Kundu S. Ann Plant Research 2000; 2(2): 161- 168.
- [13] Ramana S, Biswas AK, Kundu S, Saha JK, Yadava RBR. Bioresource Technol 2001; 82(3): 273-275.
- [14] Devarajan L, Rajanna G, Ramanathan, Oblisami G. Kisan world 1994; 21: 48-50.
- [15] Singh Y, Raj Bahadur. Indian J Agri Science 1998; 68: 70-74.
- [16] Rani R, Sri Vastava MM. Int J Ecology and Environ Science 16-23.
- [17] Rajendran K. Indian Botanical Contactor 1990; 7: 139-144.



- [18] Sahai R, Jabeen S, Saxena PK. Indian J Ecology 1983; 10: 7-10.
- [19] Chares S. Sugarcane 1985; 1: 20.
- [20] Samuel G. The use of alcohol distillery waste as a fertilizer, Proceedings of International American Sugarcane Seminar 1986; 245-252.
- [21] Chandraju S, Basavaraju HC. Sugar J (SISSTA) 2007; 20-50.
- [22] Chandaraju S, Siddappa, Chidan Kumar CS. Bio-research Bull 2007; 5:1-10.
- [23] Chandraju S, Siddappa, Chidan Kumar CS. Current Botany 2011; 3:38-42.
- [24] Manivasakam N. Physio-chemical examination of water, sewage and Industrial effluent. Pragathi Prakashan Merut 1987.
- [25] Piper CS. Soil and Plant Analysis, Han's Publication, Bombay 1966.
- [26] Jackson ML. Soil Chemical Analysis. Prentice Hall of India Pvt. Ltd., New Delhi 1973.
- [27] Walkley AJ, Black CA. Soil Sci 1934; 37: 29-38.
- [28] Subbiah BV, Asija GL. Cur Sci 1956; 25: 259-260.
- [29] Black CA. Methods of Soil Analysis. Part 2, Agronomy monograph No. 9. Am Soc Agron Madison, Wisconsin, USA 1965; 15-72.
- [30] Lindsay WL, Norve WA. Soil Sci Soc Am J 1978; 42: 421-428.
- [31] Chandraju S, Thejovathi C, Chidan Kumar CS. Plant Biol 2011; 1(2).
- [32] Chandraju S, Thejovathi C, Chidan Kumar CS. Studies on the Germination and Growth of Zinnia and Vinca seeds irrigated by Distillery spentwash. 2011; SISSTA.
- [33] Chandraju S, Thejovathi C, Chidan Kumar CS. J Chem Pharm Res 2011; 3(5):376-381.