

Research Journal of Pharmaceutical, Biological and Chemical Sciences

Safety Modern Methods for Food Using *Tetrahymena pyriformis*.

Ruslan Saferbegovich Omarov, Alexander Viktorovich Agarkov, Vladislav Evgen'evich Zakotin, and Sergei Nikolaevich Shlykov*.

Technological Management Department, Stavropol State Agrarian University, Stavropol, Russia

ABSTRACT

Over the last decade increased the number of toxicants that affect the safety of raw materials and food, simultaneously with it, time increases and methods complicated for their determination. The complete analysis of the quality and food safety for all normalized indicators is quite costly. In this regard, relevant is integrated quality control by biotesting. The purpose of work - experimental evaluation methods: determination of acute toxicity by the culture of the ciliates *Tetrahymena pyriformis*, for control the safety raw materials and food products in the process of monitoring. Objects of the study were fresh herbs (parsley, dill), wild boar, duck, beef, mincemeat (beef, pork), fish (pike). The paper presents results of studies evaluating methods by biotesting for control of food safety in the process of monitoring. In most cases, showed the correlation results of determination toxicity by methods to determining acute toxicity on ciliates, and also content specific anthropogenic and natural toxicants. The method for determining acute toxicity by ciliates is sufficiently rapid, does not require using of experimental animals or costly equipment and has prospects for acceleration of safety control raw materials and food products.

Keywords: food safety, *Tetrahymena pyriformis*, express analysis, toxicity, ciliates.

*Corresponding author

INTRODUCTION

With global food safety testing services projected to reach nearly \$20 billion by 2022, the food industry is met with increasing demands from both the consumer and regulatory agencies to maintain a safe food product.

Contaminants may occur in our food from various sources. They typically pose a health concern, resulting in strict regulations of their levels by national governments and internationally by the Codex Alimentarius Commission. Therefore, analysis of relevant chemical contaminants is an essential part of food safety testing programs to ensure consumer safety and compliance with regulatory limits. Modern analytical techniques can determine known chemical contaminants in complex food matrices at very low concentration levels. Moreover, they can also help discover and identify new or unexpected chemical contaminants.

Contaminants can be present in foods mainly as a result of the use of agrochemicals, such as residues of pesticides and veterinary drugs, contamination from environmental sources (water, air or soil pollution), cross-contamination or formation during food processing, migration from food packaging materials, presence or contamination by natural toxins or use of unapproved food additives and adulterants.

Guaranteeing safe food is a complex task. This is because food is increasingly being traded at global level, new ingredients and products are being developed, pathogens occur and consumers change their eating habits. Identifying and implementing adequate control measures is the only thing that stands between your food business producing safe food or not [1].

An important task food industry is ensuring and monitoring the safety of products. The presence in food of anthropogenic and natural toxicants is the most common cause of security breaches [2]. The former includes heavy metals, dioxins, synthetic plant growth regulators, substances used in livestock: the drug (antibiotic, antiparasitic, hormonal drugs), phosphorus and chlorogenic pesticides etc [3]. To the second group, the toxins, and metabolites of bacteria, fungi, plants [4]. Detection and identification of contaminants is not an easy task, especially if they are present at low concentration levels. It requires expertise and a good analytical strategy that is based on all gathered information about the sample and potential sources of contamination. Any clues, such as changes in smell, taste or texture, as well as a description of potential poisoning symptoms may be important in this respect. Concurrent analysis of control (“good”) samples with suspect samples is often essential to find differences and eliminate potential false positives.

At the same time, research has improved analytical methods, offering faster turnaround times for results with higher sensitivity for low-level contaminants and detection of contaminants in complex matrices. With increasing demands to test foods for more potential safety hazards, many companies look to automation. Big food producers and global food companies are trying to centralize their testing for cost savings, and this means that technology has to be set for high throughput.




Figure 1: Examples of morphotypes of Tetrahymena (a ciliate).

There are a large number of biological models for assessment toxicity, the most frequently using are laboratory animals and single-celled organisms [5]. Various modifications of the method of biotesting were

successfully used to define many objects of food production and the environment [7]. Using the *Tetrahymena pyriformis*. (figure 1) to determine the toxicity of food is seen as a promising way. The increasing use of *Tetrahymena* for both research and educational purposes has been facilitated by the ease with which it can be grown and maintained in a wide range of conditions, ranging from single cells in hanging drops to multi-liter cultures grown in large bio-reactors.

The purpose of this research is the experimental evaluation methodology for determining acute toxicity for safety control of raw materials and food in the process of monitoring.

MATERIALS AND METHODS

Objects of the study were fresh herbs (parsley, dill), wild boar, duck, beef, mincemeat (beef, pork), fish (pike). Each sample weighed approximately 500 to 550 g. Subsamples for enumeration of bacteria and flagellates were diluted and then stained with acridine orange prior to observation via epifluorescence microscopy on 0.22- μ m-pore-size Millipore black 25-mm-diameter filters, similar to the methods of Hobbie et al. [8]. Other subsamples were examined by phase-contrast microscopy under an inverted microscope. All data were converted to numbers of organisms per milliliter of the drainage from produce.

The samples were washed with sterile Tris-buffered saline solution (TBSS; 2 mM KCl, 1 mM CaCl₂, 0.5 mM MgCl₂, 1 mM Tris [pH 6.8 to 7.2]) while they were still in the closed, clean plastic bags. The samples washings were then transferred aseptically into sterile petri dishes. Autoclaved wheat grains were added to the petri dishes to enrich for protozoa in the produce wash water. Enrichment would permit easier isolation because of higher concentrations of a given organism and possibly reveal the presence of more species by inducing excystment of certain cyst-forming species. The dishes were then incubated at 25°C for 24 to 48 h. After incubation, the suspensions in the dishes were observed under an inverted microscope for the presence of protozoa.

The culture of the ciliate *Tetrahymena pyriformis* cultured into a thermostat at sterile conditions into test tubes on a nutrient medium of the following composition: bacterial peptone - 20,0; glucose - 5,0; sodium chloride - 1,0; yeast extract - 1,0 (concentration, g/dm³ distilled water). For biotesting used the 3-day culture of the ciliates. The density of suspension cells in culture was (6-8 \times 10⁴) cells/cm³. For tests used 25 to 50 g sample. Liquid products investigated in natural form or in the form of an aqueous emulsion. Solid products (25 g) had pre-crushed in a homogenizer.

For grazing experiments, 2-week-old cultures of *Glaucoma* sp. were used because of the slow growth of these ciliates, whereas 3-day-old cultures of *C. steinii* and 5-day-old cultures of the two *Tetrahymena* species were used. Ciliates were washed with TBSS by centrifugation at 500 \times g and resuspended in TBSS. Amoebae were washed from NNA plates and suspended in TBSS. The concentration of protozoa in stock suspensions was determined by differential interference contrast (DIC) microscopy. For grazing tests, 24-h-old broth cultures of *S. enterica* MB108, *E. coli* O157:H7 MB269, and *L. monocytogenes* RM2387 were washed twice by centrifugation and the final pellet was resuspended in TBSS. The concentration of bacteria in the stock suspension was determined by epifluorescence microscopy.

Cocultures of the protozoa and bacteria were established at a ratio of bacteria to protozoa of 10,000:1. Controls consisted of suspensions of washed protozoa only. Cocultures were incubated at 25°C for 24 h. All coculture experiments with bacteria and protozoa were performed twice with two replicate suspensions.

The experiment was treated as a split-plot design in time with cuts as subplots. A linear mixed model was used to estimate variance components [6]. The analysis was carried out using REML (restricted maximum likelihood) analysis in Genstat (14th edition; VSN International Ltd, Hemel Hempstead, UK).

RESULTS AND DISCUSSION

By using the methodology for determining acute toxicity for extraction toxic substances used distilled water, physiological saline or phosphate buffer (pH 7.2). After extraction, the homogenate filtered through a 2-

folded paper filter. If the pH of the investigated sample is acidic or strongly alkaline, its pH adjusted to 7.0 and 7.2 by phosphate buffer. The extraction time is 15-20 min.

Evaluation results were by results of acute toxic action tested objects on ciliates. The sample after extraction was mixed with cell culture 1:1. The method of measurement the test-response - visual assessment (counting) under the microscope number of test objects through certain intervals of time (15 min, 30 min, 1 h) (figure 2).




Figure 2: Cell of Tetrahymena strain MB1125 containing E. coli O157:H7 and an expelled vesicle outside the cell. (Left) DIC image; (right) fluorescent confocal image.

Protozoa increased in numbers after they fed on pathogens for 24 h.

Acute toxicity - the death of 100% of the ciliates within 1 h. (table 1).

Table 1: Results of control safety raw materials and foodstuff

Product type	Number of rejected samples	Method and results of safety control			
		acute toxicity on the ciliate	microbiological	high pressure liquid chromatography	Ionometric
Fresh herbs (parsley, dill)	4	Positive	Not identified	Not identified	Nitrates, nitrites
Wild boar	1	Negative	Not identified	Hydrargyrum	Not identified
Duck	1	Negative	Not identified	Plumbum	Not identified
Beef	1	Negative	Not identified	Hydrargyrum	Not identified
Mincemeat (beef, pork)	1	Positive	Salmonella	Not identified	Not identified
Fish (pike)	1	Negative	Vibriosis	Not identified	Not identified

In the monitoring process were analyzed over 1000 samples. Of them, 0.9% showed toxic effect. The results of toxicity determination by biotesting on ciliates in mostly correlated with concentration of specific toxicants parallel identified by other methods.

In the study of minced meat samples, in which by biotesting was observed the reaction of *Tetrahymena* also noted the presence of Salmonella.

CONCLUSIONS

- Data for determining the safety of various food products by biotesting acute toxicity on the ciliate, physico-chemical and microbiological methods for detection specific man-made or natural toxicants in most cases showed a correlation of results.

- The method for determining acute toxicity by ciliates is sufficiently rapid, does not require to use experimental animals or costly equipment and has prospects for accelerated safety control of raw materials and food products, however, its practical implementation should be adapted for specific objects and toxicants.

REFERENCES

- [1] Charteris W.P., Kelly P.M., Morelli L., Collins J.K. Ingredient selection criteria for probiotic microorganisms in functional dairy foods. *International Journal of Dairy Technology* 1998; 51: 123–136.
- [2] Devirgiliis C., Barile S., Perozzi, G. Antibiotic resistance determinants in the interplay between food and gut microbiota. *GEN Nutrition* 2011; 6: 275–284.
- [3] El-Shafei K., Abd El-Gawad M.A.M., Dabiza N., Sharaf O.M., Effat B.A. A mixed culture of *Propionibacterium thoenii* P-127, *Lactobacillus rhamnosus* and *Lactobacillus plantarum* as protective cultures in Kareish cheese. *Polish Journal Of Food And Nutrition Sciences* 2008; 58: 433 – 441.
- [4] Hobbie, J.E., R.J. Daley, Jasper. S. Use of Nuclepore filters for counting bacteria by fluorescence microscopy. *Applied and Environmental Microbiology* 1977; 33: 1225–1228.
- [5] Ma C.J., Wu J.M., Tsai H.L., Huang C.W., Lu C.Y., Sun L.C., Shih Y.L., Chen C.W. Prospective double-blind randomized study on the efficacy and safety of an n-3 fatty acid enriched intravenous fat emulsion in postsurgical gastric and colorectal cancer patients. *Nutrition Journal* 2015; 14: 9.
- [6] Sadovoy V., Omarov R., Shlykov S., Shchedrina T. Assessment compliance of qualitative food characteristics to standard requirements. In: 15th International Scientific Conference on Engineering for Rural Development, 25-27 May, 2016, Jelgava, Latvia 2016; 360–363.
- [7] Omarov Ruslan, Agarkov Alexander, Rastovarov Evgeny, Shlykov Sergei. Modern methods for food. In: 16th International Scientific Conference ENGINEERING FOR RURAL DEVELOPMENT, 25-27 May, 2017, Jelgava, Latvia: 960-963.
- [8] Omarov Ruslan, Gorlov Ivan, Zakotin Vladislav, Shlykov Sergei. Development of marble beef technology. In: 16th International Scientific Conference ENGINEERING FOR RURAL DEVELOPMENT, 25-27 May, 2017, Jelgava, Latvia: 956-959.