

# Research Journal of Pharmaceutical, Biological and Chemical Sciences

## Phytochemical and Antimicrobial studies of Spermacece / *Borreria Verticillata* (Rubiaceae).

Shivam Chaurasia, Piyush Yadav\*, and Shashikant Maurya.

Prasad Institute Of Technology, Department Of Pharmacy, Jaunpur – 222001, Uttar Pradesh, India.

### ABSTRACT

Borreira / Spermacece is genera of Rubiaceae sizable in tropical and subtropical America, Africa, Asia, and Europe. Based on its end result morphology they may be taken into consideration via way of means of many authors to be awesome genera and maximum others, however, choose to integrate the 2 taxa below the everyday call Spermacece. Whereas the dialogue remains unclear, on this paintings they have been taken into consideration as synonyms. Some species of those genera play an vital function in conventional medicinal drug in Africa, Asia, Europe, and South America. Some of those makes use of encompass the remedy of malaria, diarrheal and different digestive problems, pores and skin diseases, fever, hemorrhage, urinary and respiration infections, headache, irritation of eye, and gums. To date, more than 60 compounds were stated from Borreria and Spermacece species such as alkaloids, iridoids, flavonoids, terpenoids, and different compounds. Studies have showed that extracts from Borreria and Spermacece species in addition to their remoted compounds own numerous organic sports, such as anti-inflammatory, antitumor, antimicrobial, larvicidal, antioxidant, gastrointestinal, anti-ulcer, and hepatoprotective, with alkaloids and iridoids because the important energetic principles. This paper in brief critiques the ethnomedicinal makes use of, phytochemistry, and organic sports of a few remoted compounds and extracts of each genera.

**Keywords:** *Alkaloids, borreria, flavonoids, iridoids, rubiaceae, spermacece, terpenoids*

<https://doi.org/10.33887/rjpbcs/2022.13.6.1>

*\*Corresponding author*

## INTRODUCTION

The worldwide cognizance of medicinal plant life has expanded in latest years, the World Health Organization (WHO) envisioned that 80% of the populace in growing international locations depend on Antimicrobial Activity and phytochemical Screening of *Borreria verticillata* flower Bud Ethanolic Extract and Fractions conventional medication usually plant primarily based totally for number one fitness care, that is a conduct that predates civilization and it's far discovered in each society no matter its degree of improvement and sophistication. Most human beings withinside the rural regions of the sector rely in large part on herbs for remedy of numerous illnesses due to the fact they represent quintessential additives for clinical exercise and because of its low cost, smooth access, ancestral experience, safety, performance and much less facet effects Many commercially proven pills utilized in contemporary-day medication had been first of all utilized in crude shape in conventional healing practices, it's far envisioned that plant substances have furnished the fashions for 50% western pills .Plants utilized in conventional African device of medication were discovered to be lively towards a extensive kind of micro-organisms and feature continually furnished a very good source for anti-infective marketers and for healing cause to remedy numerous disorders. Recipes of vegetation are crafted from mixture of various elements of or extra vegetation or from a unmarried plant part, the elements used consist of roots, stem, stem bark, flower bud, flower, fruit, twigs, exudates and changed plant, they're ingested as decoctions, teas, infusion, pounded, soaked, squeezed and boiled, others are carried out as ointments .*Borreria verticillata*(*B. verticillata*) or *Spermacoce verticillata*, additionally called broom vassourinha plant is a perennial shrubby false-button weed herb which has huge ability as a medicinal herb It is broadly disbursed in tropical regions in Africa, Asia and America. In Nigeria, Studies have showed that extracts from exclusive elements of *B. verticillata* as nicely as its remoted compounds possesses numerous organic sports together with antibacterial, analgesic, anti-inflammatory, antitumor, antidiarrheal, larvicidal, antioxidant, gastrointestinal, antimalarial, anti-ulcer, and hepatoprotective properties and it's been generally used efficiently to treatment pores and skin diseases, toothache, diarrhea, headache, dyspepsia, leprosy, furuncles, ulcers, gonorrhoeal sores, bilharziasis and paralysis, with alkaloids and flavonoids because the major lively Therefore *B. verticillata* flower bud (aerial elements) is used as a conventional herb for its medicinal value .

### Techniques and utensils used in collection and identification of Plant substances-

The blossoming bud became accumulated in October 2019 from the Botanical nursery of the Bayero University Kano, Nigeria, and prominent at Herbarium section of the branch of Plant Biology, Bayero College Kano. The bloom bud was modified into air dried below the coloration and powdered utilizing pestle and mortar and moved into impermeable cubicles. A hundred grams (200g) of powdered example changed into permeated with ninety-six% ethanol with the aid of a sequential thorough approach (Fatope et al., 1993). The listen have become separated into boxes using a Whatman No. 1 channel paper and dissipated to dryness, gauged, and labeled. The unrefined ethanolic separate grow to be fractionated with various solvents in the growing request of extremity starting with outright n-Hexane, Chloroform and 60% Methanol as depicted through Oluwaseun et al. (2013). Antimicrobial interest and phytochemical Screening of *Borreria verticillata* blossom Bud Ethanolic Extract and Fractions

### Phytochemical screening of Extract/elements

The pay attention/divisions had been evaluated for the presence of bioactive professionals as consistent with the techniques for Arunkumar and Muthuselvam (2009) and Ushie and Adamu, (2010).

### Isolation of test Organisms

The evaluate population protected offspring of a half 12 months to 5 years who answered to the hospital with loose bowels. An easy compartment turns out to be applied for the assortment of feces exams while rectal swabs wherein accrued while the stool is not unfastened from the kids. This was completed via the researcher with the assist of a prepared lab professional. The media applied with the surrender purpose of this evaluation covered MacConkey agar, Deoxycholate agar (DCA), Blood agar, and *Salmonella Shigella* agar (SSA) and have been ready as in step with manufacturer's commands. All the examples had been delicate and brooded at 37o for 24 hours (Cheesebrough, 2010).

**Biochemical test**

A look at All plates that confirmed proof of development had been uncovered to conventional bacteriologic biochemical assessments which included: Catalase, Coagulase, Indole checks, Methyl pink test, Vegas-prosecutor take a look at, Citrate use take a look at (IMViC), Oxidase test, Urease check, sugar maturation, gasoline creation, and motility checks as portrayed by using the usage of Cheesbrough, (2010) then, at that component, progressed biochemical identity became finished using Microgen identification unit (Microbact 24E) from Oxoid Laboratory merchandise and programming adhering to the producer's instructions.

**Suspectibility test**

The agar well dispersion way have become applied as depicted through Prescott,(2002),5 (five) wells of 6mm measurement every had been made on immunized Muller Hinton agar the use of a sterile plug drill. The wells have been loaded up with various centralizations of the listen/divisions; 100mg/ml, 50mg/ml, 25mg/ml, 12.5mg/ml and DMSO (as regrettable control) and are permitted to diffuse for round 2 hours. Plates of Gentamicin have been applied as certain control. They take a look at plates that had been hatched at 37oC for 24hr. Moreover, the observed zones of the dilemma are predicted.

**RESULTS**

**Table 1: Table 1: Physical appearance and yield of extract/fractions from the flower buds of Borreria verticillata plant**

| solvent     | Extract colour | Extract texture | Weight of plant material used (gm) | Extraction yields(gm) | Percentage recovery |
|-------------|----------------|-----------------|------------------------------------|-----------------------|---------------------|
| ethanol     | brown          | Solid powder    | 200                                | 5.037                 | 2.515               |
| n-hexane    | green          | Soft powder     | 3.00                               | 0.781                 | 26.03               |
| chloroform  | green          | Soft powder     | 3.00                               | 0.654                 | 21.80               |
| 60%methanol | brown          | Soft            | 3.00                               | 0.600                 | 20.00               |

Table 1: Extraction of secondary metabolites from the flower bud of B. verticillata the usage of ninety-six% ethanol, hexane, chloroform, and 60% methanol yielded 5.037g (2.515%), zero.781g (26.03%), 0.654g (21.eighty%) and zero.60g (20.00%).

**Table 2: Preliminary phytochemicals of extract/fractions of Borreria verticillata flower bud**

| Phytochemical compound | Saponins | Pc | Glycoside | Terpenoids | Steroids | Alk | Flavnoid | RS | Tan |
|------------------------|----------|----|-----------|------------|----------|-----|----------|----|-----|
| Ethanol                | +        | +  | +         | +          | +        | +   | +        | +  | +   |
| Chloroform             | +        | -  | +         | +          | +        | +   | +        | +  | -   |
| n-hexane               | +        | -  | +         | +          | +        | +   | +        | -  | -   |
| 60% Methanol           | +        | +  | +         | +          | +        | +   | +        | +  | +   |

Table 2: It indicates the result of phytochemical screening, it is depicted that Saponins, glycoside, terpenoids, alkaloids, and flavonoids are found in all the fractions at the same time as phenolic compounds and tannins are absent within the chloroform fraction and phenolic compound, lowering sugars and tannins are absent inside the n-hexane fraction, all of the phytochemical compounds are present inside the crude ethanol and 60% methanol fractions. Those phytochemicals were stated via Mohammed et al., (2016) for antimicrobial interest with Alkaloids, tannins, flavonoids, and phenols as the

primary phytochemical ingredients. Ushie and Adamu(2010), additionally mentioned the equal phytochemicals inside the leaf extract/fractions of *B.verticillata* in which hexane, ethyl acetate, acetone, chloroform, and methanol were used in the extraction. This suggests that both the flora and leaves are similar in their phytochemical content material. This similarity is not sudden because the vegetation is changed leaves. The steroid that became detected in all the fractions, which is contrary to the work of Okwu in 2004, has been changed into observed handiest in ethyl acetate fraction of the flower bud.

#### ***Tests performed to confirm the component:***

##### **Test for alkaloids**

The extract (0.5 g) became stirred with 2 M aqueous hydrochloric acid (5.0 mL) on a steam bath. 1.0 mL of the filtrate had been one after the other dealt with with some drops of Mayer's reagent, Drangendoffs' reagent, Wagner's reagent. The ensuing answer became determined for color modifications.

##### **Test for tannins**

0.5g of every of the plant extracts became boiled with distilled water (a hundred mL) for five min. To 2.0 mL of the cooled answer (filtrate) some drops of ferric chloride became brought. The color alternate became recorded.

##### **Test for glycosides**

A small element of every of the plant extracts became positioned in separate check tubes of 0.1 M H<sub>2</sub>SO<sub>4</sub> became brought to one and distilled water (5.0 mL) brought to the other. The check tubes had been heated for forty five min in a water bath. The cooled answers had been made alkaline with an answer of 2M NaOH. Fehling answers (5.0 mL) A and B (ratio1:1) became brought to the 2 check tubes and had been allowed to face for three min. The answer of the extracts in distilled water serves as control. The modifications in response had been determined and recorded.

##### **Test for saponins**

The froth take a look at and emulsion take a look at as defined via way of means of Harborne (1975) have been used to decide the presence of saponins. A small component of every of the plant extracts changed into introduced to distilled water (20 mL) in a 100 mL beaker, boiled and filtered and the filtrate used for the take a look at; (a) Froth take a look at: five ml of the filtrate changed into diluted with water (20 mL) and shaken vigorously and allowed to face for 30 min. The end result changed into recorded. (b) Emulsion take a look at: 2 drops of olive changed into introduced to the frothing answer and shaken vigorously. The end result changed into recorded. In order to remove 'falsepositive", the blood haemolysis take a look at changed into achieved on the extract that frothed water.

##### **Test for anthraquinones**

0.5 g of every of the plant extracts changed into shaken with benzene (2.0 mL) and clear out out wherein necessary. 10 % ammonia answer (4.0 mL) changed into introduced to the filtrate. The resultant aggregate changed into shaken and the response observed and recorded.

##### **Test for flavonoids**

(a) Lead acetate take a look at: 0.5 g of the extract dissolved in 5 mL of distilled water. 10 % of lead acetate answer (1.0 mL) changed into introduced. The color formation changed into recorded. (b) Iron (III) chloride. To an answer of 0.5 g of the extract in water, drops of iron (III) chloride changed into introduced. A color extrade stated and recorded.

## Test for terpenoids

An answer of every of the extract changed into made via way of means of dissolving 0.5 g of the extract in 2.0 mL of chloroform and focused H<sub>2</sub>SO<sub>4</sub>. The presence of terpenes in the pattern changed into detected because the color changes.

## Antimicrobial Studies Of *Spermacoe Verticillata*

### Sample collection and preparation

#### Ethanollic and Hexane Extracts

Each extract changed into organized exhaustively from 50g of the dried aerial parts (BVT), roots (BVR), or the complete plant (BV). BVT1, BVR1 and BV1 have been received with the aid of using percolation with 80% ethanol to yield 2.61%, 4.32% and 3.70% w/w dry weight, respectively. BVT2, BVR2 and BV2 have been received with the aid of using percolation with hexane and the yield changed into 0.63%, 0.08% and 0.37% w/w dry weight, respectively. Each extract changed into evaporated in vacuo.

#### Preparation of Samples

10mg of every lipophilic fraction (BVT2, BVR2, BV2, BVA1 and BVA2) turned into dissolved in 0.5ml of polyethyleneglycol 400 (PEG400), then suspended in physiological tris-buffer (pH 7.4) to present a attention of 1mg/ml. Each ethanollic fraction (BVT1, BVR1, and BV1) turned into dissolved in 0.5ml of dimethylsulfoxide (DMSO), and then suspended in physiological tris-buffer to present a attention of 1mg/ml. The first terrible manipulate turned into acquired with the aid of using dissolving 0.5ml PEG400 in 9.5ml of physiological tris-buffer, and the second one terrible manipulate with the aid of using blending 0.5ml DMSO with 9.5ml physiological tris-buffer.

#### Antimicrobial Testing

The qualitative assessment of the antimicrobial pastime of the extracts turned into finished the use of the hollow plate diffusion technique for micro organism and yeasts while the strong dilution technique turned into hired for antifungal checking out the use of dermatophytes. The following microorganisms have been used : gram positive cocci which include *Staphylococcus aureus*, *Streptococcus viridans*, *Str.pneumoniae*; gram poor cocci which include *Neisseria gonorrhoeae*; acid-speedy bacilli which include *Mycobacterium fortuitum*; yeasts which include *Candida albicans* and *C. tropicalis*; the gram-variable (facultative anaerobic micro organism) *Gardnerella vaginalis*.

#### Antimicrobial Activity

The antimicrobial houses of the extracts of *B.verticillata* are summarized in table 3, and as compared with the ones of neomycin, penicillin or nystatin. The extracts of the aerial elements, the foundation and the entire plant inhibited greater or much less the increase of the examined microorganisms. But, the aerial component turned into greater lively than the entire plant which turned into additionally greater powerful than the foundation. Except for *N. gonorrhoeae*, the antibacterial impact and spectrum of the ethanol extract of the aerial elements have been greater essential than the ones of the hexane ones. *S. aureus* and *N.gonorrhoeae* have been the maximum touchy bacteria to each the ethanol and hexane extracts of the aerial elements, the entire plant or the foundation. The apolar extracts have been inactive in opposition to *Gardnerella vaginalis* and *M. fortuitum*. *Candida albicans* and *C. tropicalis* have been weakly touchy to the aerial elements and the entire plant extracts.

At a attention of 1mg/ml, the overall alkaloid extract (BVA2) became proven to own the best and widest antimicrobial properties. The antimicrobial pastime of the crude alkaloidal fraction became large towards *S. aureus*, slight towards *N. gonorrhoeae*, *Str. Pneumoniae*, *M. fortuitum*, *Gardnerella vaginalis*, *C. albicans*, *C. tropicalis*, and susceptible towards *Str. viridans*. However, those sports have been slight compared with the same old energetic compounds towards Gram-fantastic and Gram-poor cocci and acid-fast bacilli. It have to be observed that the dearth of region of inhibition does now no longer always suggest a loss of pastime for the reason that some compounds do now no longer diffuse nicely into agar.

On the opposite hand, it should be noted that the alkaloidal extracts precipitated once they have been diluted with an answer of physiological tris buffer. Such phenomenon likely diminishes the pastime. In this initial investigation, the extracts of *B. verticillata* have proven a slight antibacterial at the side of a susceptible anti-yeast sports. Such antibacterial efficiency may want to be correlated to the presence of alkaloids (BVA2) which showed the most powerful and widest spectrum antimicrobial pastime. These outcomes may want to help the conventional use of the plant in the remedy of numerous pores and skin diseases

**Table 3: Antimicrobial activity of the extracts(1mg/ml)of *B.verticilata***

| Extr. | Inhibition                   | Zone                            | Width                        | (mm)                         |                               |                                |                         |                           |
|-------|------------------------------|---------------------------------|------------------------------|------------------------------|-------------------------------|--------------------------------|-------------------------|---------------------------|
|       | <i>Staphylococcus aureus</i> | <i>Streptococcus pneumoniae</i> | <i>Neisseria gonorrhoeae</i> | <i>Gardnerella vaginalis</i> | <i>Streptococcus viridans</i> | <i>Mycobacterium fortuitum</i> | <i>Candida albicans</i> | <i>Candida tropicalis</i> |
| BVT1  | 10                           | 2                               | 12                           | 5                            | 2                             | 5                              | 3                       | 3                         |
| BVR1  | 6                            | 0                               | 10                           | 0                            | 3                             | 0                              | 0                       | 0                         |
| BV1   | 8                            | 0                               | 13                           | 5                            | 5                             | 5                              | 3                       | 3                         |
| BVT2  | 5                            | 0                               | 14                           | 0                            | 0                             | 0                              | 0                       | 0                         |
| BVR2  | 3                            | 0                               | 13                           | 0                            | 6                             | 0                              | 0                       | 0                         |
| BV2   | 5                            | 3                               | 14                           | 0                            | 5                             | 0                              | 3                       | 0                         |
| BVA1  | 5                            | 0                               | 9                            | 0                            | 0                             | 0                              | 0                       | 0                         |
| BVA2  | 16                           | 11                              | 14                           | 11                           | 9                             | 20                             | 8                       | 8                         |
| A     | 0                            | 0                               | 0                            | 0                            | 0                             | 0                              | 0                       | 0                         |
| B     | 0                            | 0                               | 0                            | 0                            | 0                             | 0                              | 0                       | 0                         |
| C     | 22                           | 20                              | 20                           |                              |                               | 40                             |                         |                           |
| D     |                              | 30                              | 30                           | 25                           | 30                            |                                |                         |                           |
| E     |                              |                                 |                              |                              |                               |                                | 20                      | 10                        |

**CONCLUSION**

The present investigation showed a conspicuous antimicrobial activity of *B. verticillata* which is closely related to the presence of alkaloids. It also showed the scientific rationale behind the traditional use of the plant in the treatment of various skin diseases.

**REFERENCES**

- [1] Abdullahi-Gero, H. S., Ahmed, A., Umar, Z. A. and Hussaini, I. M. (2014). Preliminary evaluation of ethanol flower bud extract of *Borreria verticillata* Linn (Rubiaceae) for analgesic and anti-inflammatory effects. *Journal of Medicinal Plant Research*. 8(20): 736747.
- [2] Abubakar, U. S., Yusuf, K. M., Abdu, G. T., Saidu, S. R., Jamila, G. A., and Fatima, A. (2017). Ethnopharmacological survey of medicinal plants used for the management of pediatric ailments in Kano State, Nigeria. *Research Journal of Pharmacognosy (RJP)*. 4(3): 29-39.
- [3] Adamu, H. M., Ushie, O. A. and Gwangwala, A. H. (2013). Estimation of Total Flavonoids and Tannins in the Stem Bark and Leaves of *Anogeisus leiocarpus* Plant. *Int. J. Trad. Nat. Med.* 2013, 2(3): 141-148
- [4] Anele, E.U., Amlabu, W.E. and Yashim, E.Y. (2014). Phytochemical Screening and Bacteriocidal Activity of Ethano Flower Extract of *Borreria verticillata* and *Vocanga africana*. *Nigerian Journal of Scientific Research*, 13(2): 1-3
- [5] Arunkumar, S. and Muthuselvam, M. (2009). Analysis of phytochemical constituents and antimicrobial activities of aloe vera L. against clinical pathogens. *World Journal of Agricultural Science*. 5(5): 572-576
- [6] Balde, A. M., Pietexrs, L. A., Traore, M. S., Camara, A., Balde, M. A., Oulare, K., et al., (2015). Chemotherapeutical evaluation of *Borreria verticillata* extracts. *Journal of plant Science* 3: 28-31.
- [7] Cheesebrough, M. (2010). *District Laboratory Practice in Tropical Countries part two*, New York, Cambridge University Press U.S. pp. 64-67.
- [8] Conserva, L. M. and Ferreira, J.C. (2012). *Borreria* and *Spermacoce* species (Rubiaceae): A review of their ethnomedicinal properties, chemical constituents, and biological activities. *Pharmacognosy Reviews*. 6(11): 46-55.
- [9] Fatope, M. O., Ibrahim, H. and Takeda, Y. (1993). Screening of higher plants reported as pesticides using the brine shrimp lethality assay. *International journal of phamacognosy*. 31(4): 250-56.

- [10] Mahesh, B., and Satish, S. (2008). Antimicrobial activity of some important medicinal plant against plant and human pathogens. *World Journal of Agricultural Sciences*. 4(5): 839-843.
- [11] Mohammed, A. H., Na'inna, S. Z., Yusha'u, M., Salisu, B., Adamu, U. and Garba, S. A. (2016). In vitro Assessment of Antibacterial Activity of Citrus aurantifolia Extracts. *UMYU Journal of Microbiology Research*. 1 (1): 1-6.
- [12] Newman, D. J. (2008). Natural products as leads to potential drugs: an old process or the new hope for drug discovery? *Journal of Medical Chemistry*. 51(9): 2589-2599.
- [13] Okwu, D.E. (2004) Phytochemicals and vitamin content of indigenous spices of South Eastern Nigeria. *Journal Sustainable Agriculture and Environment* 6:30-34.
- [14] Oluwaseun, M. B., Agbendeh, M. Z. and Jacob, G. A. (2013). The comparative studies of phytochemical screening of Piliostigmathonningii root and leaves extract. *Asian Journal of Plant Science and Research*. 3(6):74-77.
- [15] Prescott, M. L., Harley, J.P. & Klein, D. A. (2002). *Microbiology* (5th Ed.) McGraw-Hill Publishers, New York, USA, pp978. Shajiselvin, C.D., Kottai, M. A. and Suresh, K. (2010). Evaluation of in vivo antioxidant and lipid peroxidation effect of various extracts of the whole plant of *Borreria hispida* (Linn) on rat fed with high fat diet. *International Journal of Pharmaceutical Sciences Review and Research*. 3(1): 66-69.
- [16] Ushie, O. A. and Adamu, H. M. (2010). Phytochemical screening of *Borreria verticillata* leaves. *Journal of Agriculture, Biotechnology and Ecology*. 13(1):108-117.
- [17] Lisowski, S. (2009): *Flore (Angiospermes) de la République de Guinée. Première partie (texte)*. Jardin Botanique national de Belgique (*Scripta Botanica Belgica*, 41); 517p.
- [18] Dominique T. (1965): *Comment le Noir se soigne-t-il ? ou Médecine et Magie Africaines*. Ed., "Présence Africaine. Club Africain du Livre. [3]
- [19] Baldé A.M. (1990): *Biological and Phytochemical Investigations on Three Plants Widely Used in Guinean traditional Medicine*, PhD. Thesis, University of Antwerp, Belgium.
- [20] Ushie O. A., Adamu H. M., Ogar D. A. and Gunda H. J. (2003): Phytochemistry of *Borreria verticillata* Stem Bark. *International Journal of Traditional and Natural Medicines*. 2, 2; 97-103
- [21] Baldé A.M., A. Gergely, L.A. Pieters, M. Claeys, D.A. Vanden Berghe and A.J. Vlietinck (1989): Antimicrobial alkaloids from *Borreria verticillata*. *Planta Medica*. Supplement issue, 55 ; 652
- [22] Shajiselvin C.D., Kottai Muthu A., Suresh K.(2010) : Evaluation of in vivo antioxidant and lipid peroxidation effect of various extracts of the whole plant of *Borreria hispida* (Linn) on rat fed with high fat diet. *International Journal of Pharmaceutical Sciences Review and Research*. 3, 1 ; 66-69.
- [23] Schwartz O., Henin, Y., Marechal V. and Montagnier L. (1988): A Rapid and Simple Colorimetric Test for the Study of Anti-HIV Agents. *Aids Res. And Human Retroviruses*, 4, 6 ; 441-448.
- [24] Maynard G., Pousset J.L., Mboup S., and Denis F. (1980): Antibacterial effect of borreverine, an alkaloid isolated from *Borreria verticillata*, Rubiaceae. *C.R. Soc. Biol. (Paris)* 174 ;