

Research Journal of Pharmaceutical, Biological and Chemical Sciences

Prophylactic Antibiotic Versus Only Meticulous Pre-Operative And On The Table Proper Painting Of The Operating Site In Clean Elective Minor Surgical Cases.

Prakhar Nagar¹, Vijay Pal¹, Shwetabh Pradhan^{2*}, and Avinash Sharma².

¹Assistant Professor, Department OF General Surgery, JNUIMSCRC, Jaipur, Rajasthan, India.

²Associate Professor, Department OF General Surgery, GMC Haldwani, Rajasthan, India.

ABSTRACT

Preoperative day, night and operative day, on the table meticulous skin preparation of the operating site. This is done with povidone-iodine scrub-paint combination (Betadine) with an isopropyl alcohol application. This gives comparable results in preventing infection. Thus avoids unnecessary exposure of antibiotics and cost effectiveness to the patient. Keywords: Antibiotic abuse, Skin preparation, Prophylactic antibiotic.

Keywords: Antibiotic abuse, Skin preparation, Prophylactic antibiotic.

<https://doi.org/10.33887/rjpbcs/2023.14.6.4>

**Corresponding author*

INTRODUCTION

This is a study conducted in the clean elective minor case to study about antibiotics versus skin preparation with betadine and ethyl alcohol. Recently it was been noticed that antibiotics abuse is being common in the modern era. Hence for clean elective case just the good skin preparation on table is more efficient and essential to prevent infection and antibiotic abuse was been prevented. In this study a 100 case was being selected and being studied.

Aim And Objectives

Preoperative day, night and operative day, on the table meticulous skin preparation of the operating site. This is done with povidone-iodine scrub-paint combination (Betadine) with an isopropyl alcohol application.

This gives comparable results in preventing infection. Thus avoids unnecessary exposure of antibiotics and cost effectiveness to the patient.

METHODS

Inclusion Criteria

Clean elective cases eg: Inguinal, epigastric, umbilical Hernias, Solitary nodule thyroid, Hydrocele, Circumcision and appendicectomy.

Age up to 30 years(Thirty years). Cases operated GMC HALDWANI

Exclusion Criteria

Co-morbid diseases, like diabetic, hypertension, skin diseases, anemia.

Complicated hernias. Proved and doubtful thyroid malignancies. Circumcision cases with cracked prepuces, paraphimosis.

Methodology

Povidone-iodine scrub-paint combination (Betadine) with an isopropyl alcohol application is used for skin preparation.

Thorough skin preparation was done on the night of pre-operative day.

Preparation done from nipple to knee in all hernias and appedecectomies, umbilicus to knee in circumcision, hydrocele and front of neck shoulder up to mid-upper arm in thyroid.

Same team will do the same skin preparation on the table before draping.

Study period one year 2022 to 2023.

Pilot study of 70 cases (varying samples in all cases)

DISSCUSION

In our hospital settings both private and Government, usage of prophylactic antibiotic is administrated as a matter of routine practice with few exceptions. Dosage, time and duration differs.

Method I: One full dose of 3rd generation cephalosporins on pre-operative day night, one dose on the table-peroperatively and one dose on that day night.

Method II: One full dose on the table-peroperatively and one dose on that day night.



Method III: Broad spectrum antibiotic parenterally (Amoxicillin 500mg BID or 3rd generation cephalosporins OD dose for first three days and same drug oral till sutured removed (varies from Five to seven days).

Why prophylactic antibiotics required in our settings and same not required in western country settings.

Our setting environment is not very concussive for preventing infection. Weather humid and ours is a tropical country.

Whereas western country settings environment is clean and contusive due to their weather condition. Prophylactic antibiotics may not be required as a matter of routine.

Present study disproves the above notions. Most of the clean elective minor surgeries, infection goes from with out to with in than inherent infection with in. Hence through skin preparation is sufficient and any method of prophylactic antibiotic is not required.

Post-operative infection rate is marginally less in cases with out prophylactic antibiotics, which proves skin preparation is more important than antibiotics. Medical economics, wastage of money. Exposure of patients to drugs with out indication.

Patients has to suffer from pain of IV cannula. Added risk of hospital acquired infection if cannula is kept for more number of days. Additional burden of work and wastage of man-hours of work of Interns and paramedics which may hamper their quality of work.

Unnecessary hospital stay affecting monetary wise to the health care provider and associated expenditure for the patient, loss wages on over stay in hospital. Loss of work force in patients office / work place [1-5].

CONCLUSION

This pilot study indicates the following facts

Antibiotic abuse on elective clean selected cases.

Loss of money to the health provider both direct and indirect.

Loss of money to the patient both direct and indirect.

Quality of work by both medical and paramedical staff.

Hence a meta-analysis or cohorts randomized control study to be done with a larger sample.

ACKNOWLEDGMENTS

The Study conducted on in-patients of Govt Medical college Haldwani, thanks to all the Assistant professors. Residents, Interns and Paramedics.

REFERENCES

- [1] Guidelines for Antimicrobial Usage 2008-2009 Tables 29 - 31 from the Cleveland Clinic
- [2] Disease Management Project Online Medical Reference from the Cleveland Clinic
- [3] Page C P, Bohnen J M, Fletcher J R, McManus A T, Solomkin J S, Wittmann D H. Antimicrobial prophylaxis for surgical wounds. Guidelines for clinical care. Arch Surg 1993;128:79-88.
- [4] Khera SY, Kostyal DA, Deshmukh N. A comparison of chlorhexidine and povidone-iodine skin preparation for surgical operations. Curr Surg 1999;56:341-343.
- [5] Armstrong EP, Patrick KL, Erstad BL. Comparison of preoperative skin preparation products. Pharmacotherapy 2001;21:345-350.